

Global and uniqueness properties of stationary and static spacetimes with outer trapped surfaces

MARC MARS

Facultad de Ciencias, Universidad de Salamanca

marc@usal.es

MARTIN REIRIS

Albert Einstein Institut, Max Planck Institut für Gravitationsphysik

martin@aei.mpg.de

Global properties of maximal future Cauchy developments of stationary, m -dimensional asymptotically flat initial data with an outer trapped boundary are analyzed. We prove that, whenever the matter model is well posed and satisfies the null energy condition, the future Cauchy development of the data is a black hole spacetime. More specifically, we show that the future Killing development of the exterior of a sufficiently large sphere in the initial data set can be isometrically embedded in the maximal Cauchy development of the data. In the static setting we prove, by working directly on the initial data set, that all Killing prehorizons are embedded whenever the initial data set has an outer trapped boundary and satisfies the null energy condition. By combining both results we prove a uniqueness theorem for static initial data sets with outer trapped boundary.

Contents

1	Introduction and overview of the main results	2
2	Stationary Killing initial data	4
2.1	Background and definitions	4
2.2	The statements of the main results: Theorems 4 and 5	8
2.3	Structure and proof of Theorem 4.	9
2.4	Proofs of the auxiliary Propositions 2, 3, 4, 5 and of the Theorem 5.	11
3	Static Killing initial data	20
3.1	Background and definitions	20
3.2	The statements of the main results: Theorem 6 and Corollary 1	22
3.3	Volume monotonicity along “optic” congruences of geodesics	22
3.4	On the volume of horizons of asymptotically flat static KIDs	26
4	Uniqueness of static, vacuum, asymptotically flat initial data sets with outer trapped boundary	32
5	Acknowledgments	34
6	Appendix A: Fixed points have $I_1 < 0$	35

1 Introduction and overview of the main results

In this paper we investigate the relationship between asymptotically flat stationary initial data sets with outer trapped boundary and black holes. By Penrose singularity theorem [18] initial data configurations with an outer trapped boundary lead to maximal Cauchy developments which are geodesically incomplete. On the other hand, physical arguments of predictability lead to the weak cosmic censorship conjecture (see e.g. [22]) which asserts that any singularity that forms during a process of gravitational collapse must be a black hole space-time. Black hole space-times (from now on simply *black holes*) satisfy strong global properties (see below) and it is a very difficult task in general to determine whether the global properties will be satisfied knowing only the initial configuration of the spacetime. In the stationary setting, where to a certain sense there is no evolution at all, the problem must necessarily be much simpler. At first sight one might even think that, in fact, determining whether the maximal Cauchy development of a stationary initial configuration with trapped boundary is a black hole should be direct because one would only need to propagate the initial information by the isometry. The problem is not nearly as simple because even being able to carry over the development by the isometry no global information, for example global hyperbolicity, is a priori evident.

There exist at least two different approaches to show that the maximal globally hyperbolic development of a stationary initial data set is a black hole. The first approach tries to prove directly that the maximal globally hyperbolic development of the data enjoys sufficient global properties to qualify as a black hole. This is the approach we follow in the first part (Section 2) of this paper.

There exist several definitions of black hole, a priori non equivalent, but following the same underlying principle. In a stationary setting, a convenient definition requires the existence of an asymptotically flat $(m + 1)$ -end with the property that its causal past does not cover the whole space-time manifold. An asymptotically flat $(m + 1)$ -end is the natural generalization to higher dimensions of the usual notion of asymptotically flat four-end, which can be found e.g. in [1]. In essence, the definition demands that the $(m + 1)$ -end is topologically the product of a real line times the exterior of a closed ball in \mathbb{R}^m . The spacetime metric is required to be constant along the \mathbb{R} factor and to approach the Minkowski metric (with time along the \mathbb{R} factor) at suitable rate. Since we are interested only in the future of a Cauchy surface, we restrict the topology of the $(m + 1)$ -end to be \mathbb{R}^+ times the exterior of a closed ball in \mathbb{R}^m . Under suitable conditions, this definition is equivalent to the definition using conformal compactifications (see e.g. the Appendix of [15] for the vacuum, four dimensional case or Proposition 1.9 in [10] for electrovacuum).

Concerning stationary initial data and black holes we will prove in Section 2

Theorem 1 *Let \mathcal{D} be an m -dimensional ($m \geq 3$) asymptotically flat stationary initial data with well posed matter model satisfying the null energy condition. Suppose that $\partial\Sigma \neq \emptyset$ is future outer trapped. Then the maximal Cauchy development (\mathcal{M}, g) of \mathcal{D} is a black hole*

spacetime.

A more precise statement is given in Theorem 4 in Section 2.2, where we prove that the Killing development of the data outside a sufficiently large sphere can be isometrically embedded into the maximal globally hyperbolic spacetime generated by the whole data.

The second approach to prove that equilibrium initial configurations lead to black holes is via uniqueness theorems. In the stationary setting, black hole space-times satisfy uniqueness theorems, in particular for arbitrary dimension in the static case and in dimension four in the non-static electrovacuum case under suitable hypotheses [12]. Thus, if one expects that certain stationary initial data develops well behaved black-hole space-times then such data should be embeddable in one of the stationary/static black holes allowed. In other words the data (inside some region) should be one among those data endowed on sections of the listed black holes. The hope is that, somehow, such information should be extractable from the initial data itself to deduce, a fortiori, that the given initial data gives rise to a black-hole space-time.

This strategy has been successfully applied in the past under suitable restrictions. The first result along these lines is due to P. Miao [20] who proved a uniqueness result for asymptotically flat, three-dimensional vacuum and time-symmetric static Killing initial data having an outermost minimal boundary. More precisely, Miao proved that the data must be isometric to the $\{t = \text{const.}\}$ slice of the Schwarzschild spacetime for some mass $M > 0$. A related result was found by Carrasco & Mars [5, 6] for data with outer trapped boundary in the case of non-zero second fundamental form and for more general matter models, provided a number of conditions were satisfied. The generalization to non-vanishing second fundamental form is relevant because, in the absence of global information about the spacetime generated by the initial data, globally defined time-symmetric slices may simply not exist in the spacetime under consideration. Although of interest, the results in [5, 6] are not fully satisfactory because they required a number of hypotheses that were basically dictated by the method of proof, with no fundamental reason to believe that they should be necessary. One such hypothesis excluded the presence of so-called non-embedded Killing prehorizons. A Killing prehorizon is a null immersed hypersurface where the Killing vector is null, nowhere zero and tangent (see [12]). When the surface gravity of the horizon vanishes, it is a priori possible that the Killing prehorizon is not embedded (see the discussion in the Addendum of [9]). In a black hole context (more precisely, assuming that the domain of outer communications is globally hyperbolic) Chruściel and Galloway [8] have proved that all prehorizons contained in the domain of outer communications must be embedded. Therefore, in the light of the discussion above, one expects to be able to rule out non-embedded prehorizons at the initial data level.

Our second aim in this paper, developed in Section 3, is precisely to exclude the existence of non-embedded Killing prehorizons in the exterior region of a static Killing initial data set. Note that since our global statement in the first part of the paper is only to the future, we cannot rely on the results by Chruściel and Galloway mentioned above. This has the advantage that not even the existence of a spacetime containing the initial data needs to be

assumed. This means, in particular, that no field equations whatsoever are required for this part of the work. The only requirement we make is that the matter model satisfies the null energy condition. In this part of the paper, however, we restrict ourselves to static Killing initial data. It is an interesting open question whether the method extends to the stationary (non-static) setting as well.

Concerning the relation between static Killing initial data and horizons we will be proving in Section 3

Theorem 2 *Let \mathcal{D} be an m -dimensional ($m \geq 3$) asymptotically flat static Killing initial data set satisfying the null energy condition. Suppose that the exterior, connected region where the Killing vector is timelike lies in the interior of Σ , then each degenerate horizon is a compact embedded submanifold.*

Combined with Theorem 1, this result implies the non-existence of non-embedded Killing prehorizons in static, asymptotically flat spacetimes with Cauchy surface having an outer trapped boundary.

The third, and final, part of the paper, developed in Section 4, is an application of the previous two and establishes a uniqueness theorem for static, asymptotically flat initial data with outer trapped boundary for suitable matter models. In the vacuum case, the statement is as follows

Theorem 3 *Let \mathcal{D} be a static, vacuum m -dimensional ($m \geq 3$) asymptotically flat Killing initial data with non-empty future outer trapped boundary. Then, the initial data restricted to the exterior, connected region where the Killing vector is timelike can be isometrically embedded in a Schwarzschild $(m + 1)$ -dimensional spacetime of mass $M > 0$.*

The non-vacuum case is treated in Theorem 7 in Section 4. This statement gives a satisfactory answer to the problem of uniqueness for static initial data sets with an outer trapped boundary.

2 Stationary Killing initial data

2.1 Background and definitions

In this paper manifolds are defined to be smooth, Hausdorff, connected and paracompact (hence second countable). Fields on manifolds are assumed to be smooth. For manifolds with boundary Σ we use $\partial\Sigma$ for the boundary and $\Sigma^\circ (= \Sigma \setminus \partial\Sigma)$ for the usual notion of interior of a manifold with boundary. For arbitrary subsets U in a manifold we use \overline{U} for the topological closure, $\text{Int}(U)$ for the interior and $\partial^T U$ for the topological boundary.

We work with $(m + 1)$ -dimensional spacetimes (\mathcal{M}, g) ($m \geq 3$). For a subset $U \subset \mathcal{M}$ we define the *null boundary* $\partial^N U$ as the subset of points $p \in \partial^T U$ such that there exists a future directed null geodesic segment $\gamma(s)$ of (\mathcal{M}, g) starting at p and fully contained in $\partial^T U$.

The causal past of a set U is denoted by $J^-(U)$ and the future domain of dependence of U is denoted by $D^+(U)$. The conventions we use for these objects, and for causality notions in general, follow [25]. In particular, a spacetime (\mathcal{M}, g) is globally hyperbolic if it admits a Cauchy hypersurface Σ . We denote by \mathcal{M}^+ the future domain of dependence of Σ° . Note that \mathcal{M}^+ is a manifold with the smooth boundary Σ° . Let n be the space-time future directed unit normal to Σ° . The induced metric on Σ° will be denoted by g and the second fundamental form by K (in the direction of n). If Σ has boundary we assume it is compact and that both g and K extend smoothly to $\partial\Sigma$.

Let \mathbf{G} be the Einstein tensor of g , namely $\mathbf{G} := \text{Ric} - \frac{1}{2}\mathbf{R}g$, where Ric denotes the Ricci tensor of the metric g and \mathbf{R} is the curvature scalar (our sign conventions are such that the Ricci tensor and curvature scalar of a round sphere are positive)

We define ρ , J (1-form) and \mathcal{T} (symmetric two-tensor) at $p \in \Sigma$ according to

$$\begin{aligned} \rho &= \mathbf{G}(n, n), \\ (1) \quad J(v) &= -\mathbf{G}(n, v), \quad v \in T_p\Sigma, \\ \mathcal{T}(v, w) &= \mathbf{G}(v, w), \quad v, w \in T_p\Sigma. \end{aligned}$$

Remark 1 ρ, J, \mathcal{T} are defined in terms of the Einstein tensor, and not as components of any energy-momentum tensor because we will not assume any specific field equations relating the Einstein tensor with the energy-momentum tensor of the matter fields. Neither the matter model nor the field equations will be of concern to us except that we will require a well posedness property defined later.

The data $(\Sigma; (g, K); (\rho, J))$ satisfies the energy and momentum constraint equations

$$\begin{aligned} (2) \quad R_g - |K|_g^2 + k^2 &= 2\rho, \\ (3) \quad \text{div}_g (K - kg) &= -J. \end{aligned}$$

where $k = \text{tr}_g K$ and div_g is the divergence with respect to the metric g .

The boundary of Σ (if any) is said to be *future outer trapped* if it is compact and

$$\theta^+(\partial\Sigma) := \text{tr}|_{\partial\Sigma} K + H < 0.$$

where $\text{tr}|_{\partial\Sigma} K$ is the trace of K restricted to $\partial\Sigma$ and H is the mean curvature of $\partial\Sigma$ in the inward direction to Σ .

We will assume that we have a Killing vector field ξ on (\mathcal{M}, g) having the decomposition $\xi = Nn + Y$ along Σ . (N, Y) will be part of the initial data. Because ξ is Killing we have, over Σ , the equations (see e.g. [13])

$$\begin{aligned} (4) \quad \mathcal{L}_Y g &= -2NK, \\ (5) \quad \mathcal{L}_Y K &= -\text{Hess}_g N + N(\text{Ric}_g + kK - 2K \circ K) - N\left(\mathcal{T} - \frac{1}{m-1}(\text{tr}_g \mathcal{T} - \rho)g\right). \end{aligned}$$

where \mathcal{L} denotes Lie derivative, Hess_g is the Hessian with the metric g , Ric_g is the Ricci tensor of g and $K \circ K$ is the tensor obtained from $K \otimes K$ by tracing with g the second and fourth indices. Concerning the data we make the following definition.

Definition 1 (AF-KID) *The data $\mathcal{D} := (\Sigma; (g, K); (\rho, J, \mathcal{T}); (N, Y))$ is said to be a*

1. *Killing initial data (KID) if it satisfies (2)-(3) and (4)-(5),*
2. *and stationary asymptotically flat (AF) if there is a compact set whose complement Σ^∞ is diffeomorphic to $\mathbb{R}^{m+1} \setminus \{\text{closed ball}\}$ and, in the Euclidean coordinates $\bar{x} = (x^1, \dots, x^{m+1})$ on Σ^∞ defined by the diffeomorphism we have*

$$(6) \quad g_{ij} - \delta_{ij} = O^2(1/r^{m-2}), \quad K_{ij} = O^2(1/r^{m-1}),$$

$$(7) \quad N - N_\infty = O^2(1/r^{m-2}), \quad Y_i - Y_{\infty i} = O^2(1/r^{m-2}).$$

and with $r = |\bar{x}| := \sqrt{(x^1)^2 + \dots + (x^m)^2}$, where N_∞, Y_∞ are constants satisfying $N_\infty > |Y_\infty|$.

Under these conditions we will say simply that \mathcal{D} is a stationary asymptotically flat initial data.

Let $\lambda := N^2 - |Y|_g^2$. Since $\lambda \rightarrow N_\infty^2 - |Y_\infty|^2 > 0$ at infinity, we can assume (after restricting Σ^∞ if necessary) that $\lambda > 0$ on the asymptotically flat end Σ^∞ . We denote by Σ^T (T from time-like) the connected component of $\{\lambda > 0\} \subset \Sigma$ containing Σ^∞ .

The Killing development of subregions of Σ^T is defined as follows.

Definition 2 (Killing developments) *Let Ω be a connected open subset of Σ^T . Then the infinite Killing development of the data at Ω is defined as the space-time*

$$(8) \quad \mathcal{K}(\Omega) := (\Omega \times (0, \infty), g_D = -\lambda dt^2 + Y \otimes dt + dt \otimes Y + g),$$

where $Y := g(Y, \cdot)$. The restriction of $\mathcal{K}(\Omega)$ to $t \in (0, \bar{t}]$ ($\bar{t} > 0$) is denoted by $\mathcal{K}(\Omega, \bar{t})$.

The Killing development is a space-time with Killing field $\xi = \partial_t$ and Einstein tensor \mathbf{G} as defined above from ρ, J and \mathcal{T} .

In this paper we will also use a related notion of Killing development of hypersurfaces V (with or without boundary) embedded in a spacetime (\mathcal{M}, g) admitting a Killing vector ξ . The only requirement is that V is everywhere transverse to ξ . If we denote by g the first fundamental form of V , by Y the pull-back of the one-form obtained by lowering the index to ξ and $\lambda := -\langle \xi, \xi \rangle$ (where $\langle \cdot, \cdot \rangle$ denotes scalar product with the spacetime metric g) then, the Killing development $\mathcal{K}(V)$ is defined as the spacetime $V \times (0, \infty)$ with the metric g_D defined exactly as in (8). It is immediate to see that, ξ being transverse to V everywhere, g_D is a metric of Lorentzian signature. There is no restriction on the causal character of V , which

in particular is allowed to be null. Note that when the Killing initial data is embedded in a spacetime, we have a priori two ways of constructing the Killing development. However, the spacetimes constructed with either definition are the same, so no confusion arises. We also emphasize that the Killing development $\mathcal{K}(V)$ is an abstract spacetime defined on its own which, a priori, has nothing to do with the original spacetime (\mathcal{M}, g) .

Given (\mathcal{M}, g) with a Killing vector ξ , let $\beta_q(\lambda)$, $\lambda \geq 0$, be the Killing orbit starting at $q \in \mathcal{M}$ (i.e. $\beta_q(\lambda = 0) = q$). For any $W \subset \mathcal{M}$ and $0 \leq a \leq b$ satisfying the property that all the Killing orbits $\beta_q(\lambda)$, $q \in W$ extend to all values $\lambda \in [a, b]$ we will denote by $O_{[a,b]}(W)$ the set

$$O_{[a,b]}(W) := \{\beta_q(\lambda) \in \mathcal{M} / q \in W, a \leq \lambda \leq b\}.$$

If $a = b$ we will simply write $O_{[a]}(W)$ for $O_{[a,a]}(W)$. In the following it will be convenient to use two different notations for Killing orbits: $\beta_q(\lambda)$ when the orbit satisfies $\beta_q(\lambda = 0) = q$ and λ takes positive values and $\alpha_p(\mu)$ when the orbit satisfies $\alpha_p(\mu = 0) = p$ and μ takes negative values.

Definition 3 (Well posed matter models) *Let \mathcal{D} be a Killing initial data. We say that the matter model is well posed if the field equations are such that a Killing initial data generates a unique maximal globally hyperbolic space-time (\mathcal{M}, g) having a Killing field ξ extending $\xi|_{\Sigma} = Nn + Y$.*

As usual, we will refer to the maximal globally hyperbolic spacetime (\mathcal{M}, g) as *maximal Cauchy development*. The simplest example of well posed matter model is vacuum, i.e. when the field equations are $\mathbf{G} = 0$. In this case, the Cauchy problem is well posed and the maximal Cauchy development admits a Killing vector [21]. The same is true e.g. in electrovacuum [10] and for many other matter models [24].

We will say that a matter model satisfies the null energy condition if all Cauchy developments (\mathcal{M}, g) solving the field equations satisfy $\mathbf{G}(l, l) \geq 0$ for any null vector l .

We need some observations on isometric embeddings. Let $(\mathcal{N}_i, g_{\mathcal{N}_i})$, $i = 1, 2$ be two connected Lorentzian or Riemannian manifolds (possibly with smooth boundary). Let $\mathcal{W}_i \subset \mathcal{N}_i^\circ$, $i = 1, 2$ be two sets such that $\mathcal{W}_i \subset \overline{\text{Int}(\mathcal{W}_i)}$, $i = 1, 2$ and $\text{Int}(\mathcal{W}_i)$, $i = 1, 2$ are connected and non-empty. Let $p_i \in \text{Int}(\mathcal{W}_i)$, $i = 1, 2$. Let $\phi : T_{p_1}\mathcal{N}_1 \rightarrow T_{p_2}\mathcal{N}_2$ be a linear isometry between the tangent space to \mathcal{N}_1 at p_1 and the tangent space to \mathcal{N}_2 at $p_2 = \phi(p_1)$. Then we will write $(\mathcal{N}_1, \mathcal{W}_1) \Subset_\phi (\mathcal{N}_2, \mathcal{W}_2)$ if there is an open set U_1 of \mathcal{N}_1 containing \mathcal{W}_1 and a smooth map $\varphi : U_1 \rightarrow \mathcal{N}_2$ such that $\varphi|_{\text{Int}(\mathcal{W}_1)}$ is an isometric embedding of $\text{Int}(\mathcal{W}_1)$ into $\text{Int}(\mathcal{W}_2)$. In such case $\varphi|_{\mathcal{W}_1}$ is unique (and determined only by ϕ). In the rest of the work we will simply write $\mathcal{W}_1 \Subset \mathcal{W}_2$ and it should be understood that this also entails the existence of companion manifolds $\mathcal{N}_1, \mathcal{N}_2$ and map ϕ . The companion objects will be understood from the context. Note the transitivity property: $\mathcal{W}_1 \Subset \mathcal{W}_2, \mathcal{W}_2 \Subset \mathcal{W}_3 \Rightarrow \mathcal{W}_1 \Subset \mathcal{W}_3$. When $\mathcal{W}_1 \Subset \mathcal{W}_2$ we will say that \mathcal{W}_1 lies in \mathcal{W}_2 .

The uniqueness of the maximal future globally hyperbolic space-time \mathcal{M}^+ implies that

any future globally hyperbolic space-time $(\mathcal{N}, g_{\mathcal{N}})$ with Cauchy hypersurface $C = \partial\mathcal{N}$ isometric to a particular connected open subset of (Σ, g) , lies uniquely inside (\mathcal{M}^+, g) . In other words $\mathcal{N} \in \mathcal{M}^+$. This fact will be of fundamental importance.

We will denote by S_r the coordinate sphere of coordinate radius r in the asymptotically flat end Σ^∞ . S_r separates Σ into two closed parts (i.e. including their boundaries), $\Sigma^E(r)$ (“E” from “Exterior”) and $\Sigma^I(r)$ (“I” from “Interior”). Note that $S_r \subset \Sigma^T$ and $\Sigma^E(r) \subset \Sigma^T$. For $d \geq 0$, we also define $T(\partial\Sigma, d) := \{p \in \Sigma / \text{dist}_g(p, \partial\Sigma) \leq d\}$ and $\Sigma_d := \{p \in \Sigma / \text{dist}_g(p, \partial\Sigma) \geq d\}$.

Definition 4 *Let (\mathcal{M}, g) be a globally hyperbolic space-time with Cauchy surface Σ and asymptotically flat stationary data \mathcal{D} . Suppose $\mathcal{K}(\Sigma^E(r)) \Subset \mathcal{M}^+$. Then we define the future event horizon (over Σ) as the topological boundary of $J^-(\mathcal{K}(\Sigma^E(r))) \cap \Sigma$ (as a subset of Σ).*

The Killing development $\mathcal{K}(\Sigma^E(r))$ is a future asymptotically flat $(m+1)$ -end as described in the Introduction.

2.2 The statements of the main results: Theorems 4 and 5

The following theorem is a precise version of Theorem 1 in the Introduction.

Theorem 4 *Let \mathcal{D} be an asymptotically flat, stationary initial data set with well posed matter model satisfying the null energy condition. Suppose that $\partial\Sigma$ (if non-empty) is future outer trapped. Then the maximal Cauchy development (\mathcal{M}, g) of \mathcal{D} satisfies*

1. *There is $r > 0$ such that $\mathcal{K}(\Sigma^E(r))$ lies in \mathcal{M}^+ .*
2. *There is $d > 0$, such that $T(\partial\Sigma, d) \cap J^-(\mathcal{K}(\Sigma^E(r))) = \emptyset$.*

In basic terms, if the boundary of Σ is future outer trapped, then the future event horizon over Σ exists, is computable or constructible from the data and does not intersect $\partial\Sigma$. Therefore $\partial\Sigma$ lies inside the future black hole region.

The constant r in item 1 is introduced in Proposition 1 later. The constant d is any positive constant satisfying the property that $\partial\Sigma_{d'}, 0 \leq d' \leq 2d$, is diffeomorphic to $\partial\Sigma$ and $\theta^+(\partial\Sigma_{d'}) < 0$.

We also have

Theorem 5 *Let \mathcal{D} be an asymptotically flat stationary initial data with well posed matter model satisfying the null energy condition. Suppose that $\partial\Sigma \neq \emptyset$ is future outer trapped. Then, the exterior of the event horizon contains Σ^T . In particular, $\Sigma^T \cap \partial\Sigma = \emptyset$.*

2.3 Structure and proof of Theorem 4.

The proof is constructive. We define first the future globally hyperbolic region E_0^+ . The definition will come out from the following proposition which we leave without proof.

Proposition 1 *Consider the domains of dependence $D^+(\Sigma^E(r))$ depending on r , as sets inside the Killing development $\mathcal{K}(\Sigma^E(r))$. Then, if r is big enough the null boundary $\partial^N D^+(\Sigma^E(r))$ is a smooth null hypersurface foliated by future complete null geodesic rays starting at S_r . Moreover $D^+(\Sigma^E(r)) \Subset \mathcal{M}^+$.*

Fixed one such r we will denote from now on $E_0^+ := D^+(\Sigma^E(r))$ and its Cauchy surface $\Omega_0 = \Sigma^E(r)$. Now, starting from E_0^+ , we will construct inductively a sequence $E_0^+ \Subset E_1^+ \Subset E_2^+ \Subset \dots \Subset E_\infty^+ = \bigcup_{i \geq 0} E_i^+$, such that $E_i^+ \setminus \partial^N E_i^+$ are globally hyperbolic spacetimes with Cauchy surfaces $\Omega_0 \Subset \Omega_1 \Subset \Omega_2 \Subset \dots \Subset \Omega_\infty := \bigcup_{i \geq 0} \Omega_i \Subset \Sigma^\circ$ none touching the tubular neighborhood $T(\partial\Sigma, d)$ for some $d > 0$, namely having $\Omega_\infty \cap T(\partial\Sigma, d) = \emptyset$. From the uniqueness of the maximally globally hyperbolic development \mathcal{M}^+ we conclude that $E_i^+ \Subset \mathcal{M}^+$ for all i and therefore that $E_\infty^+ \Subset \mathcal{M}^+$. Moreover we show explicitly along the construction that $\mathcal{K}(\Sigma^E(r)) \Subset E_\infty^+$. This, together with $E_\infty^+ \Subset \mathcal{M}^+$, gives $\mathcal{K}(\Sigma^E(r)) \Subset \mathcal{M}^+$ which is the claim (1) in Theorem 4. On the other hand $J^-(\mathcal{K}(\Sigma^E(r))) \cap T(\partial\Sigma, d) \subset E_\infty^+ \cap T(\partial\Sigma, d) = \Omega_\infty \cap T(\partial\Sigma, d) = \emptyset$ which is the claim (2), and last, in Theorem 4. We present now progressively the main definitions and Propositions (auxiliary Propositions 2-5) leading to the construction of the sequence $\{E_i\}$. The construction, together with the proof of Theorem 4, is explained after the statement of Proposition 5 which is the main statement of this section and which in itself structures the inductive procedure.

An important collection of regions for the proof are \mathcal{W}_t , $t \geq 0$ defined as

$$(9) \quad \mathcal{W}_t := \mathcal{K}(\partial^N E_0^+, t) \cup E_0^+ \subset (\Sigma^E(r) \times [0, \infty), g_D).$$

Note that \mathcal{W}_t is a Lorentzian manifold with smooth boundary and corners. Suppose that $\mathcal{W}_t \Subset \mathcal{M}^+$, $t \geq 0$. We consider the sets

$$\begin{aligned} E_t &:= J^-(\mathcal{W}_t), \\ E_t^+ &:= E_t \cap \mathcal{M}^+, \end{aligned}$$

where here and in the following J^- is taken in the spacetime (\mathcal{M}, g) . When $t = 0$ then $E_t^+ = E_0^+$ and $\partial^N E_0^+$ is, as we said, smooth. Moreover we will prove

Proposition 2 *ξ points strictly outwards from E_0^+ at $\partial^N E_0^+$.*

Crucially, this property is generalizable to the sets E_t , $t \geq 0$ which could fail to have smooth boundaries (although their boundaries are Lipschitz manifolds). We prove that ξ points strictly outwards from E_t at $\partial^N E_t$ in the sense of the following definition which generalizes the standard sense that we understood in Proposition 2.

Definition 5 Suppose that $\mathcal{W}_t \in \mathcal{M}^+$. Let $E_t = J^-(\mathcal{W}_t)$. Then, we say that ξ points strictly outwards to E_t at $\partial^N E_t$, if for every $p \in \partial^N E_t$, there is $\mu_p < 0$ such that

D1-1. $\alpha_p(\mu) \in \text{Int}(E_t)$, for all μ with $\mu_p \leq \mu < 0$, where $\alpha_p(\mu)$ is the Killing orbit passing through p at $\mu = 0$,

D1-2. there is an m -manifold $V_p \subset \text{Int}(E_t)$, transversal to ξ at $\alpha_p(\mu_p)$ such that, if we denote by $\beta_q(\lambda)$, $\lambda \geq 0$ the Killing orbit passing through $q \in V_p$ at $\lambda = 0$ then

D1-2-(a). there is a first $\lambda > 0$, denoted by λ_q , for which $\beta_q(\lambda_q) \in \partial^N E_t$,

D1-2-(b). the map $q \rightarrow \beta_q(\lambda_q)$ from V_p into $\partial^N E_t$ is continuous

D1-2-(c). for every $q \in V_p$, $\beta_q(\lambda) \in (\mathcal{M} \setminus E_t)$ if $\lambda > \lambda_q$ but close to it,

In other words ξ points strictly outwards to E_t at its boundary if every Killing orbit starting at $\text{Int}(E_t)$ either remains inside $\text{Int}(E_t)$ or crosses $\partial^N E_t$. An obvious consequence of the Definition is that if an orbit starts in $\text{Int}(E_t)$ and crosses $\partial^N E_t$, then it never returns to E_t .

Recalling, we will prove

Proposition 3 Suppose that $\mathcal{W}_t \in \mathcal{M}^+$. Let $E_t = J^-(\mathcal{W}_t)$. Then ξ points strictly outwards from E_t at $\partial^N E_t$ in the sense of Definition 5.

On the other hand if ξ points strictly outwards from E_t at $\partial^N E_t$ in the sense of Definition 5, then we prove that one can abstractly extend E_t “along the Killing” any time $\bar{t} > 0$ in the sense of the following definition which generalizes the notion of Killing development introduced in Definition 2 before.

Definition 6 Suppose that $\mathcal{W}_t \in \mathcal{M}^+$. Let $E_t = J^-(\mathcal{W}_t)$. The infinite abstract Killing development of E_t , $K(E_t)$, is defined as the manifold formed by the open sets $\{\text{Int}(E_t), \mathcal{K}(V_p), p \in \partial^N E_t\}$ endowed with their respective metrics and subject to the following identifications

D2-1. the points $x_1 = (q, \lambda) \in \mathcal{K}(V_p) = V_p \times (0, \infty)$ and $x_2 \in \text{Int}(E_t)$ are identified iff $\lambda < \lambda_q$ and $\beta_q(\lambda) = x_2$.

D2-2. the points $x_1 = (q_1, \lambda_1) \in \mathcal{K}(V_{p_1})$ and $x_2 = (q_2, \lambda_2) \in \mathcal{K}(V_{p_2})$ are identified iff

D2-2-(a). $\beta_{q_1}(\lambda_{q_1}) = \beta_{q_2}(\lambda_{q_2})$ and,

D2-2-(b). $\lambda_{q_1} - \lambda_1 = \lambda_{q_2} - \lambda_2$.

The abstract Killing development at time $\bar{t} \geq 0$ is defined as the following region of $K(E_t)$

$$K(E_t, \bar{t}) := \text{Int}(E_t) \cup \{\beta_q(\lambda + \lambda_q), q \in V_p, p \in \partial^N E_t, 0 \leq \lambda \leq \bar{t}\}.$$

Similarly one defines

$$K(E_t^+, \bar{t}) := (\text{Int}(E_t^+) \cap \mathcal{M}^+) \cup \{\beta_q(\lambda + \lambda_q), q \in V_p, p \in \partial^N E_t^+, 0 \leq \lambda \leq \bar{t}\}.$$

Of course we have $E_t^+ \Subset E_t \Subset K(E_t, \bar{t})$. However one must think $K(E_t, \bar{t})$ as a new spacetime bearing a priori no global relation with \mathcal{M} . In general one would not expect that $K(E_t, \bar{t}) \Subset \mathcal{M}$. As we will explain in Proposition 5 the situation will be different for $K(E_t^+, \bar{t})$ if we select \bar{t} properly and this is what will allow us eventually to construct the sequence $\{E_i^+\}$.

Recalling, we will prove

Proposition 4 *Suppose that $\mathcal{W}_t \Subset \mathcal{M}^+$. Let $E_t = J^-(\mathcal{W}_t)$. Then, for any $\bar{t} \geq 0$ the abstract Killing development $K(E_t, \bar{t})$ is a smooth, Lorentzian and second countable manifold with null and Lipschitz boundary.*

The following proposition will structure the construction of the sequence $\{E_i^+\}$ that we explain thereafter.

Proposition 5 *There is $t_* > 0$ depending only on the initial data over $\Sigma_{d/2} \cap \Sigma^I(r+1)$ such that if for some $t \geq 0$, we have*

$$\mathbf{H1.} \quad \mathcal{W}_t \Subset \mathcal{M}^+,$$

$$\mathbf{H2.} \quad T(\partial\Sigma, 2d) \cap E_t^+ = \emptyset \text{ with } E_t^+ := J^-(\mathcal{W}_t) \cap \mathcal{M}^+,$$

then,

$$\mathbf{C1.} \quad K(E_t^+, t_*) \Subset \mathcal{M}, \text{ and therefore } \mathcal{W}_{t+t_*} \Subset \mathcal{M}^+.$$

$$\mathbf{C2.} \quad T(\partial\Sigma, 2d) \cap J^-(K(E_t^+, t_*)) = \emptyset.$$

$$\mathbf{C3.} \quad E_{t+t_*}^+ := J^-(\mathcal{W}_{t+t_*}) \cap \mathcal{M}^+ = J^-(K(E_t^+, t_*)) \cap \mathcal{M}^+$$

We are ready to prove Theorem 4.

Proof of Theorem 4: To construct the sequence $\{E_i^+\}$ we proceed as follows. First, when $i = 0$, E_0^+ is as we defined it before. Now, **H1** and **H2** hold in Proposition 5 with $t = 0$. Then, conclusion **C1** gives $\mathcal{W}_{t_*} \Subset \mathcal{M}^+$ which is **H1** with $t = t_*$. In addition conclusions **C2** and **C3** give $E_{t_*}^+ \cap T(\Sigma, 2d) = \emptyset$ with $E_{t_*}^+ = J^-(\mathcal{W}_{t_*}) \cap \mathcal{M}^+$ which is **H2** with $t = t_*$. Then define $E_1^+ = J^-(\mathcal{W}_{t_*}) \cap \mathcal{M}^+$. Applying repeatedly Proposition 5 in this way, we are led to define E_i^+ as $E_i^+ = J^-(\mathcal{W}_{it_*}) \cap \mathcal{M}^+$ which is the desired sequence. \square

2.4 Proofs of the auxiliary Propositions 2, 3, 4, 5 and of the Theorem 5.

Proof of Proposition 2: The Proposition is direct from the fact that, as a field inside $\mathcal{K}(\Sigma^E(r))$, ξ is time-like and future-pointing and that $\partial^N E_0^+$ is smooth and null. An alternative argument (to be used later) comes from the observation that on a smooth null boundary like $\partial^N E_0^+$, proving that ξ points strictly outwards is equivalent to prove that for any null geodesic $\gamma(\tau)$ in $\partial^N E_0^+$ (parametrized by affine parameter τ into the future direction) we have $\langle \gamma', \xi \rangle < 0$. As ξ is a Killing field we have

$$\langle \gamma'(\tau), \xi(\gamma(\tau)) \rangle = \langle \gamma'(0), \xi(\gamma(0)) \rangle,$$

where $\gamma(0)$ is the initial point of the geodesic at S_r . As ξ is timelike on S_r it is $\langle \gamma'(0), \xi(\gamma(0)) \rangle < 0$. The statement follows. \square

The following lemma is useful for the proof of Proposition 3.

Lemma 1 *Assume that $\mathcal{W}_t \in \mathcal{M}^+$ and let $E_t = J^-(\mathcal{W}_t)$. Then, the closure of $\text{Int}(E_t) \cap \partial^N \mathcal{W}_t$ is compact. Moreover, any null geodesic γ in $\partial^N E_t$ satisfies $\langle \gamma', \xi \rangle < 0$.*

Proof: Assume that $\text{Int}(E_t) \cap \partial^N \mathcal{W}_t$ is not compact. Then there is a divergent sequence of points $\{q_i\}$, $q_i \in \partial^N \mathcal{W}_t$, a sequence of points $\{p_i\} \in \mathcal{W}_t$, and a sequence of timelike and past directed curves $\Gamma_i(\tau)$, $\tau \in [0, 1]$ (in \mathcal{M}^+), such that, for every $i \geq 0$, Γ_i starts at p_i and ends at q_i . However, as Γ_i is timelike and past directed, we claim that to reach q_i , Γ_i must first cross the set $U = \{\beta_q(\lambda), q \in \partial\Omega_0, 0 \leq \lambda \leq t\}$. Indeed to reach q_i from its future the curve Γ_i must first leave \mathcal{W}_t , but being timelike and past directed it cannot cross $\partial^N \mathcal{W}_t$, nor it can enter E_0^+ for it could not leave E_0^+ again. The claim follows. Denote by $\text{Vol}(\Gamma_i)(\tau) = \text{Volume}_g(J^-(\Gamma_i(\tau)) \cap \mathcal{M}^+)$. Then $\text{Vol}(\Gamma_i)(\tau)$ is, for every i , a monotonically decreasing function of τ . Moreover as $\{q_i\}$ is a divergent sequence we must have $\text{Volume}_g(J^-(q_i) \cap \mathcal{M}^+) = \text{Vol}(\Gamma_i)(1) \rightarrow \infty$. On the other hand as, for every i , Γ_i crosses U , and $\text{Vol}(\Gamma_i)(\tau)$ is monotonically decreasing, it must be $\text{Vol}(J^-(\Gamma_i))(1) \leq \sup\{\text{Vol}(J^-(q) \cap \mathcal{M}^+), q \in U\} < \infty$ for all $i \geq 0$, which gives a contradiction. An important consequence of this, to be used later, is that every inextensible future null geodesic γ in $\partial^N E_t$ becomes eventually a null geodesic of $\partial^N \mathcal{W}_t$. Constancy of $\langle \gamma', \xi \rangle$ along this geodesic proves the claim $\langle \gamma', \xi \rangle < 0$. \square

Proof of Proposition 3: We show first **D1-2-(a)**. Through every point p in $\partial^N E_t$ there passes a future inextensible null geodesic $\gamma_p(\tau)$, $\tau > 0$, starting at p and fully contained in $\partial^N E_t$ (see [25]). By Lemma 1 $\langle \gamma', \xi \rangle < 0$. Moreover, every point $p' = \gamma_p(\tau)$, $\tau > 0$, is a smooth point of $\partial^N E_t$. But for smooth points we know that if $\langle \gamma'_p(\tau), \xi(p') \rangle < 0$ then ξ points strictly outwards to E_t at p' . If $\tau > 0$ is small enough, then $p \in \partial J^-(\gamma_p(\tau))$ and p is a smooth point of $\partial^N J^-(\gamma_p(\tau))$. But because $\langle \gamma'_p(0), \xi(p) \rangle = \langle \gamma'_p(\tau), \xi(\gamma_p(\tau)) \rangle < 0$, we deduce that $\xi(p)$ points strictly outwards to $J^-(\gamma_p(\tau)) \subset E_t$ at p . Since $\text{Int}(J^-(\gamma_p(\tau)) \subset \text{Int}(E_t)$ there is $\mu_p < 0$ such that $\alpha_p(\mu) \in \text{Int}(E_t)$ for $\mu \in [\mu_p, 0)$ thus showing **D1-1**.

We prove now **D1-2-(a)**. First note that there is $a > 0$ and a closed smooth three-submanifold V_p (with smooth boundary) transversal to ξ and containing $\alpha_p(\mu_p)$ such that for every $q \in V_p$ the Killing orbit $\beta_q(\lambda)$, passing through q at $\lambda = 0$, extends to all values $\lambda \in [0, -\mu_p + a]$. For every $q \in V_p$ define $\bar{\lambda}_q = \inf\{a, \lambda_q^T\}$ where λ_q^T is the first $\lambda > 0$ such that $\beta_q(\lambda) \in \partial^N E_t$ (we take $\lambda^T = \infty$ if $\beta_q(\lambda)$ never intersects $\partial^N E_t$). Note that if $\bar{\lambda}_q < a$ then $\beta_q(\bar{\lambda}_q) \in \partial^N E_t$. From this and because $\bar{\lambda}_{\alpha_p(\mu_p)} = -\mu_p$ we deduce that if $q \rightarrow \bar{\lambda}_q$ is continuous at $q = \alpha_p(\mu_p)$ then one can take a smaller V_p if necessary in such a way that $\bar{\lambda}_q < a$ for every $q \in V_p$ and therefore with $\beta_q(\bar{\lambda}_q) \in \partial^N E_t$ as claimed in **D1-1-(a)**. We prove now the continuity of $\bar{\lambda}_q$ at $\bar{q} = \alpha_p(\mu_p)$ ($\bar{q} = \alpha_p(\mu_p)$ from now on).

First, from the proof of **D1-1** one knows that p is a smooth point of the boundary of a past cone $J^-(\gamma(\tau))$ entirely included in E_t . Moreover ξ points strictly outwards to $J^-(\gamma(\tau))$

at p . It follows that for any sequence $q_j \rightarrow \bar{q}$ we have $\liminf\{\bar{\lambda}_{q_j}\} \geq \bar{\lambda}_{\bar{q}}$. Indeed, let $\tilde{\lambda}_{q_j}$ be the first $\lambda > 0$ the orbit reaches the smooth boundary of $J^-(\gamma(\tau))$ near p . Since $\tilde{\lambda}_{q_j} \rightarrow -\mu_p$ and $\bar{\lambda}_{q_j} \geq \tilde{\lambda}_{q_j}$ the claim follows. We need to prove therefore that $\limsup\{\bar{\lambda}_{q_j}\} \leq \bar{\lambda}_{\bar{q}}$. Suppose instead that there is a sequence $q_j \rightarrow \bar{q}$ such that $\limsup\{\bar{\lambda}_{q_j}\} > \bar{\lambda}_{\bar{q}} + b$, for some $b > 0$ and $b < a$. Since E_t is closed (this is proved easily), it follows that the piece of orbit $\{\beta_{\bar{q}}(\lambda), 0 \leq \lambda \leq \bar{\lambda}_{\bar{q}} + b\}$ lies inside E_t . We claim that, as a consequence, there are points $\beta_{\bar{q}}(\lambda)$, for $\lambda > \bar{\lambda}_{\bar{q}}$ but arbitrarily close to $\bar{\lambda}_{\bar{q}}$ lying in the interior of E_t . If not, we would have that for $\lambda > \bar{\lambda}_{\bar{q}}$, the points $\beta_{\bar{q}}(\lambda)$ must lie in $\partial^N E_t$. But by **D1-1**, if a point in an orbit belongs to $\partial^N E_t$, then the points (in the orbit) near it and in the direction opposite to ξ are interior points to E_t , which is a contradiction to the fact $\beta_{\bar{q}}(\bar{\lambda}_{\bar{q}}) \in \partial^N E_t$. Thus, the orbit $\beta_{\bar{q}}(\lambda)$ satisfies the following properties:

1. $\beta_{\bar{q}}(\bar{\lambda}_{\bar{q}}) \in \partial^N E_t$,
2. $\beta_{\bar{q}}(\lambda) \in \text{Int}(E_t)$, for $0 \leq \lambda < \bar{\lambda}_{\bar{q}}$,
3. there are points $\beta_{\bar{q}}(\lambda) \in \text{Int}(E_t)$, for $\lambda > \bar{\lambda}_{\bar{q}}$ but arbitrarily close to it,

Let us show that these three facts together contradict **D1-1**. We work now with the notation $\alpha_p(\mu) = \beta_{\bar{q}}(\bar{\lambda}_{\bar{q}} + \mu)$ instead of the notation $\beta_{\bar{q}}(\lambda)$. Let $\mu_1 > 0$ be such that $\alpha_p(\mu_1)$ belongs to the interior E_t . Let now $\gamma(s)$, $s \geq 0$ be a future directed time-like geodesic, starting at p . Consider the orbits $\alpha_{\gamma(s)}(\mu)$, with $\alpha_{\gamma(s)}(0) = \gamma(s)$ and $s > 0$, but close to it. We observe that $\gamma(s) \notin E_t$ (otherwise $p \in \text{Int}(E_t)$) and if $s > 0$ is small enough then $\alpha_{\gamma(s)}(\mu_1)$ belongs to the interior of E_t . Thus we have

$$\alpha_{\gamma(s)}(\mu_1) \in \text{Int}(E_t), \quad \alpha_{\gamma(s)}(0) \in (\mathcal{M}^+ \setminus E_t).$$

Since $\mu_1 > 0$ is as small as desired this immediately contradicts **D1-1** and **D1-2-(a)** is proved.

Thus the map $q \rightarrow \beta_q(\bar{\lambda}_q)$ (making V_p smaller if necessary) is from V_p into $\partial^N E_t$. We have then $\lambda_q = \bar{\lambda}_q$ for λ_q as defined in Definition 5. Now, the argument that showed the continuity of $\bar{\lambda}_q$ at $q = \bar{q}$ shows the continuity of λ_q at any point $q \neq \bar{q}$, namely **D1-2-(b)**, and also **D1-2-(c)**. \square

Proof of Proposition 4: The fact that the infinite Killing development is a smooth manifold is seen as follows. The transition functions from $\mathcal{K}(V_p)$ into $\text{Int}(E_t)$ (on their domains of identification) according to identification **D2-1** are trivially diffeomorphisms because they are given by

$$(q, \lambda) \rightarrow \beta_q(\lambda),$$

for $q \in V_p$ and $0 < \lambda < \lambda_q$. Consider now the transitions functions from $\mathcal{K}(V_{p_1})$ into $\mathcal{K}(V_{p_2})$ (on their domains of identification) according to the identifications **D2-2**. We show that they are also diffeomorphisms. First we show that the transition functions are one to one and then we show that they are locally differentiable. Suppose that $x_1 = (q_1, \lambda_1)$ and $x'_1 = (q'_1, \lambda'_1)$

in $\mathcal{K}(V_{p_1})$ are identified to $x_2 = (q_2, \lambda_2)$ in $\mathcal{K}(V_{p_2})$ via **D2-2**. Then, because $\beta_{q_1}(\lambda_{q_1})$ and $\beta_{q'_1}(\lambda_{q'_1})$ must both be equal to $\beta_{q_2}(\lambda_{q_2})$ it follows $q_1 = q'_1$ and $\lambda_{q_1} = \lambda_{q'_1}$. On the other hand

$$\lambda_{q_1} - \lambda_1 = \lambda_{q_2} - \lambda_2, \text{ and } \lambda_{q'_1} - \lambda'_1 = \lambda_{q_2} - \lambda_2$$

Thus $\lambda_1 = \lambda'_1$ and therefore $x_1 = x'_1$. This shows that the transition functions from $\mathcal{K}(V_{p_1})$ into $\mathcal{K}(V_{p_2})$ (on their domains of identifications) are one to one. We show now that they are locally differentiable. Suppose that $x_1 = (q_1, \lambda_1) \in \mathcal{K}(V_{p_1})$ and $x_2 = (q_2, \lambda_2) \in \mathcal{K}(V_{p_2})$, are identified according to **D2-2**. Then, we have $\beta_{q_1}(\lambda_{q_1}) = \beta_{q_2}(\lambda_{q_2})$. Let V be a smooth three-manifold (without boundary) transversal to ξ everywhere satisfying $V \subset \mathcal{K}(V_{p_1}) \cap \text{Int}(E_t)$, $V \subset \mathcal{K}(V_{p_2}) \cap \text{Int}(E_t)$ (both intersections under the natural identification **D2-1**) and such that the orbit $\beta_{q_1}(\lambda)$ intersects V . Let

$$\hat{\varphi}_1 : B_1 \subset V_{p_1} \rightarrow \mathcal{K}(V_{p_1}),$$

be the embedding satisfying $\hat{\varphi}_1(B_1) = V$. It is clear that B_1 is an open neighbourhood of V_{p_1} around q_1 . As a simple example V could be chosen as the image of the graph

$$\begin{aligned} \hat{\varphi}_1 : B_1 \subset V_{p_1} &\rightarrow \mathcal{K}(V_{p_1}), \\ \bar{q}_1 &\rightarrow \hat{\varphi}_1(\bar{q}_1) = (\bar{q}_1, \lambda_{q_1} - \epsilon) \end{aligned}$$

where B_1 is a sufficiently small open neighborhood of V_{p_1} around q_1 and $\epsilon > 0$ is a sufficiently small fixed number, both chosen in such a way that $V \equiv \hat{\varphi}_1(B_1) \subset \mathcal{K}(V_{p_2}) \cap \text{Int}(E_t)$

Since $V \subset \mathcal{K}(V_{p_2})$, there exists a neighbourhood B_2 of q_2 in V_{p_2} and an embedding $\hat{\varphi}_2 : B_2 \rightarrow \mathcal{K}(V_{p_2})$ such that $\hat{\varphi}_2(B_2) = V \subset \mathcal{K}(V_{p_2})$. Restricting $\hat{\varphi}_1, \hat{\varphi}_2$ to their images, we have two diffeomorphisms

$$\varphi_1 : B_1 \rightarrow V, \quad \varphi_2 : B_2 \rightarrow V.$$

Consider now two open sets \tilde{B}_1 and \tilde{B}_2 defined by

$$\begin{aligned} \tilde{B}_1 &= \{(\bar{q}_1, \lambda), \bar{q}_1 \in B_1, \\ &\quad \lambda \in (\lambda(\hat{\varphi}_1(\bar{q}_1)) - \lambda(\hat{\varphi}_1(q_1)) + \lambda_1 - \delta, \lambda(\hat{\varphi}_1(\bar{q}_1)) - \lambda(\hat{\varphi}_1(q_1)) + \lambda_1 + \delta)\}, \\ \tilde{B}_2 &= \{(\bar{q}_2, \lambda), \bar{q}_2 \in B_2, \\ &\quad \lambda \in (\lambda(\hat{\varphi}_2(\bar{q}_2)) - \lambda(\hat{\varphi}_2(q_2)) + \lambda_2 - \delta, \lambda(\hat{\varphi}_2(\bar{q}_2)) - \lambda(\hat{\varphi}_2(q_2)) + \lambda_2 + \delta)\} \end{aligned}$$

where $\delta > 0$ is chosen sufficiently small so that $\tilde{B}_1 \subset \mathcal{K}(V_{p_1})$ and $\tilde{B}_2 \subset \mathcal{K}(V_{p_2})$. The map

$$\phi : \tilde{B}_1 \rightarrow \tilde{B}_2,$$

defined by $\phi(\bar{q}_1, \lambda) = (\varphi_2^{-1} \circ \varphi_1(\bar{q}_1), \lambda - \lambda(\hat{\varphi}_1(\bar{q}_1)) + \lambda(\hat{\varphi}_2(\varphi_2^{-1}(\varphi_1(\bar{q}_1))))$ is the transition function according to **D2-2** restricted to \tilde{B}_1 and is a smooth diffeomorphism onto its image

\tilde{B}_2 .

The Hausdorff property of the abstract Killing development is seen as follows. If $x_1 = (q_1, \lambda_1) \in \mathcal{K}(V_{p_1})$ and $x_2 \in \text{Int}(E_t)$ are different points then either $\lambda_1 \geq \lambda_{q_1}$ or not, but if not then $\beta_{q_1}(\lambda_1) \neq x_2$. In either case it is straightforward to find separating neighborhoods. Now, if $x_1 = (q_1, \lambda_1) \in \mathcal{K}(V_{p_1})$ and $x_2 = (q_2, \lambda_2) \in \mathcal{K}(V_{p_2})$ and different points then either $\beta_{q_1}(\lambda_{q_1}) \neq \beta_{q_2}(\lambda_{q_2})$ or not, and if not then $\lambda_1 \neq \lambda_2$. Also in any of these possibilities it is straightforward to find separating neighborhoods.

To see that the abstract Killing development is second countable use that $\partial^N E_t$ is a Liptschitz three-manifold, pick a dense and countable set of points $\{p_i\}$ in $\partial^N E_t$ and over each point find a V_{p_i} and construct $\mathcal{K}(V_{p_i})$. Finally define the countable open subsets of $\mathcal{K}(V_{p_i})$, $U_{ijklm} = B_{ij} \times (k/l - 1/m, k/l + 1/m)$ (k, l naturals and $k/l > 1/m$) where B_{ij} is a countable basis of open sets of V_{p_i} . The sets $\{U_{ijklm}\}$ together with a countable basis of open sets of $\text{Int}(E_t)$ gives a countable basis for the abstract Killing development. \square

Before going into the proof of Proposition 5 we need to introduce some sets and their terminology. They are in fact simple regions of \mathcal{M} although their precise definitions are somehow lengthy. The relevant sets to be used in the proof of the Proposition 5 are: the slabs $D_i, i = -3, \dots, 3$, the layers $L_i, i = 1, 2$ and the bands $B_i^j, i = 2, 3$. The graphic representation of the sets can be seen in Figure 1.




Figure 1: Schematic figure illustrating the definitions of *slabs* $D_{-3}, D_{-2}, D_{-1}, D_1, D_2, D_3$, *layers* L_{-2}, L_2 and *bands* B_1^2, B_1^3 . The initial Cauchy surface Σ , the modified Cauchy surface Σ_τ and the sets E_0^+ and E_t are also shown. The set D used in the text is the union of all regions in grey in the figure.

Given $p \in \Sigma^\circ$ we consider the space-time (inextensible) timelike geodesic $\gamma_p(\tau) \subset \mathcal{M}$ starting perpendicularly to p into the future and parametrized by proper time τ . Now, given

Ω a compact region in Σ° , we define the tubular neighborhoods

$$U(\Omega; \bar{\tau}_1, \bar{\tau}_2) := \{\gamma_p(\tau), p \in \Omega, \bar{\tau}_1 \leq \tau \leq \bar{\tau}_2\}$$

We define also

$$\begin{aligned}\bar{\partial}U(\Omega; \bar{\tau}_1, \bar{\tau}_2) &:= \{\gamma_p(\tau), p \in \Omega, \tau = \bar{\tau}_2\}, \\ \underline{\partial}U(\Omega; \bar{\tau}_1, \bar{\tau}_2) &:= \{\gamma_p(\tau), p \in \Omega, \tau = \bar{\tau}_1\}\end{aligned}$$

For $\bar{\tau}$ small enough these are compact sets inside \mathcal{M} . We fix now $\tilde{\tau} > 0$ such that

- P1.** For every $p \in \Sigma_{d/2} \cap \Sigma^I(r+1)$, the geodesic $\gamma_p(\tau)$ is defined at least on the interval $[-3\tilde{\tau}, 3\tilde{\tau}]$. Moreover the map from $(\Sigma_{d/2} \cap \Sigma^I(r+1)) \times [-3\tilde{\tau}, 3\tilde{\tau}]$ into \mathcal{M} , given by $(p, \tau) \rightarrow \gamma_p(\tau)$, is a diffeomorphism into the image.
- P2.** There is a Cauchy surface $\underline{\Sigma}_{\tilde{\tau}}$ for \mathcal{M} containing the boundary $\underline{\partial}U(\Sigma_{d/2} \cap \Sigma^E(r+1); -3\tilde{\tau}, 3\tilde{\tau})$ and coinciding with Σ outside a compact set in Σ° .
- P3.** The sets $U(\partial\Sigma_{d/2}; -3\tilde{\tau}, 3\tilde{\tau})$ and $U(S_{r+1}; -3\tilde{\tau}, 3\tilde{\tau})$ do not intersect the set

$$\left(D^+(\Sigma_d) \cup J^-(\Sigma_d)\right) \setminus E_0^+$$

Then, given such $\tilde{\tau}$, we define the *slabs*

$$\begin{aligned}D_i &:= \left(D^+(\Sigma_d) \cap U(\Sigma_{d/2} \cap \Sigma^I(r+1); 0, i\tilde{\tau})\right) \setminus E_0^+, \quad i = 1, 2, 3, \\ D_i &:= \left(J^-(\Sigma_d) \cap U(\Sigma_{d/2} \cap \Sigma^I(r+1); i\tilde{\tau}, 0)\right) \setminus J^-(E_0^+), \quad i = -1, -2, -3, \\ D &:= D_{-3} \cup D_3.\end{aligned}$$

Define the *layers* L_2 and L_{-2} as

$$\begin{aligned}L_2 &= \left(D^+(\Sigma_{2d}) \cap \bar{\partial}U(\Sigma_{d/2} \cap \Sigma^I(r+1); 0, 2\tilde{\tau})\right) \setminus E_0^+, \\ L_{-2} &= \left(J^-(\Sigma_{2d}) \cap \underline{\partial}U(\Sigma_{d/2} \cap \Sigma^I(r+1); -2\tilde{\tau}, 0)\right) \setminus J^-(E_0^+)\end{aligned}$$

Finally, suppose that $\mathcal{W}_t \Subset \mathcal{M}^+$ and let $E_t = J^-(\mathcal{W}_t)$. Then, define the bands B_t^i , $i = 2, 3$ and their upper and lower boundaries $\bar{\partial}B_t^i, \underline{\partial}B_t^i$

$$\begin{aligned}B_t^i &= (\partial^N E_t) \cap U(\Sigma_{d/2} \cap \Sigma^I(r+1); -i\tilde{\tau}, i\tilde{\tau}), \\ \bar{\partial}B_t^i &= B_t^i \cap \bar{\partial}U(\Sigma_{d/2} \cap \Sigma^I(r+1), -i\tilde{\tau}, i\tilde{\tau}), \\ \underline{\partial}B_t^i &= B_t^i \cap \underline{\partial}U(\Sigma_{d/2} \cap \Sigma^I(r+1); -i\tilde{\tau}, i\tilde{\tau}) \\ B_t^{i\circ} &= B_t^i \setminus (\bar{\partial}B_t^i \cup \underline{\partial}B_t^i)\end{aligned}$$

Proof of Proposition 5: We define t_* as the supremum of the times $\bar{t} > 0$ such that every Killing orbit $\beta_q(\lambda)$, $0 \leq \lambda < 2\bar{t}$, where $q \in L_{-2} \cup L_2$, and $\beta_q(0) = q$ lies inside $\text{Int}(D_3 \setminus D_1) \cup \text{Int}(D_{-3} \setminus D_{-1})$. Note that we are taking the range of λ between 0 and $2\bar{t}$ and not between 0 and \bar{t} .

We proceed now with the proof. Assume then **H1** and **H2**. We prove first **C1**. We note two important observations concerning Killing orbits starting at B_t^2 that will be relevant for the discussion that follows.

O-1. For any $p \in B_t^2$ the Killing orbit $\beta_p(\lambda)$, $\lambda > 0$ remains inside $\text{Int}(D \setminus E_t)$ until a first λ when it reaches $\partial^T(D \setminus E_t) \setminus B_t^3$. (The orbit cannot touch B_t^3 because ξ points strictly inwards to $D \setminus E_t$ at B_t^3).

O-2. Because of **O-1**, every pair of orbits $\beta_{p_1}(\lambda)$, $\lambda \in (0, \lambda_1)$ and $\beta_{p_2}(\lambda)$, $\lambda \in (0, \lambda_2)$ lying in $\text{Int}(D \setminus E_t)$, with $p_1, p_2 \in B_t^2$ but different, do not intersect.

We prove now that $O_{[0, 2t_*]}(B_t^2) \subset (D \setminus \text{Int}(E_t))$. Assume that such is not the case and let \bar{t}_m be the minimum of the times \bar{t} , with $0 < \bar{t} < t_*$ and such that

$$O_{[0, 2\bar{t}]}(B_t^2) \subset \mathcal{M} \text{ and } O_{[2\bar{t}]}(B_t^2) \cap \left(\mathcal{M} \setminus (D \setminus \text{Int}(E_t)) \right) \neq \emptyset$$

Let $\bar{p} \in B_t^2$ be such that

$$p = O_{[2\bar{t}_m]}(\bar{p}) \in \left(\partial^T(D \setminus \text{Int}(E_t)) \setminus B_t^3 \right)$$

where we are assuming that p is not in B_t^3 because of **O-1**. Let $\gamma(\tau)$, $\tau \in [0, 1]$, be a past directed null geodesic inside B_t^2 starting at $\bar{q} \in \bar{\partial}B_t^2$ and ending at \bar{p} . Then $O_{[2\bar{t}_m]}(\gamma(\tau))$ is a past directed null geodesic starting at $q = O_{[2\bar{t}_m]}(\bar{q})$ and ending at p . But by definition of \bar{t}_m , it is $\bar{t}_m < t_*$ and therefore it must be $q \in \text{Int}(D_{-3} \cup D_3)$. Because of **O-1** the geodesic $O_{[2\bar{t}_m]}(\gamma(\tau))$ cannot intersect B_t^3 . Therefore by the definition of D it must be $p \in \Sigma_{\bar{t}}$. That this is an impossibility is seen as follows. First note that

$$\partial^T O_{[0, 2\bar{t}_m]}(B_t^2) = B_t^{2\circ} \cup O_{[2\bar{t}_m]}(B_t^{2\circ}) \cup O_{[0, 2\bar{t}_m]}(\underline{\partial}B_t^2) \cup O_{[0, 2\bar{t}_m]}(\bar{\partial}B_t^2)$$

and that because of **O-2** the union on the right hand side is disjoint. Second we claim that inextensible past directed time-like geodesics $\Gamma(\tau)$, $\tau \geq 0$, starting at the point p (found before) at $\tau = 0$ must remain inside $\text{Int}(O_{[0, 2\bar{t}_m]}(B_t^2))$ (for $\tau > 0$) until a first $\tau = \bar{\tau}$ when it reaches

$$(10) \quad B_t^{2\circ} \cup O_{[0, 2\bar{t}_m]}(\underline{\partial}B_t^2) \cup O_{[0, 2\bar{t}_m]}(\bar{\partial}B_t^2)$$

Indeed if instead there is such a $\Gamma(\tau)$ and $\bar{\tau} > 0$ with $\Gamma(\bar{\tau}) \in O_{[2\bar{t}_m]}(B_t^{2\circ})$ then $O_{[-2\bar{t}_m]}(\Gamma(\tau))$, with τ near $\bar{\tau}$, would be a past directed time-like geodesic inside E_t and crossing $B_t^{2\circ}$ at $O_{[-2\bar{t}_m]}(\Gamma(\bar{\tau}))$ which is not possible as $B_t^2 \subset \partial^N E_t = \partial^N J^-(\mathcal{W}_t)$. Thus any past directed

time-like geodesic Γ starting at p would eventually touch (10). But because the set (10) lies to the future of $\underline{\Sigma}_{\bar{t}}$ and $p \in \underline{\Sigma}_{\bar{t}}$ we obtain an impossibility. We have thus proved that $O_{[0,2t_*]}(B_t^2) \subset (D \setminus \text{Int}(E_t))$ as we wanted.

Because of this and because of **O-2** we claim that we can construct a natural differentiable map from $O_{(0,2t_*)}(B_t^{2\circ})$ into $K(E_t, 2t_*)^\circ$, which is actually an isometry. In other words we claim that we have naturally $O_{(0,2t_*)}(B_t^{2\circ}) \subseteq K(E_t, 2t_*)^\circ$. Roughly speaking the isometry can be explained in the following terms: We can think of B_t^2 both as a set in \mathcal{M} or as a set in $K(E_t, 2t_*)$ then the map identifies Killing orbits in \mathcal{M} starting at points in $B_t^{2\circ}$, as a set in \mathcal{M} , with Killing orbits in $K(E_t, 2t_*)$ starting in $B_t^{2\circ}$, but now as a set inside $K(E_t, 2t_*)$. In precise terms, the map is defined as follows. Let o be a point in $O_{(0,2t_*)}(B_t^{2\circ})$. We will define the map in a neighborhood of it. We can write $o = O_{[\bar{t}]}(p)$ with $0 < \bar{t} < 2t_*$, and with $p \in B_t^{2\circ}$. Both, \bar{t} and p , are unique because of **O-2**. Let $\mu_p < 0$, $q = \alpha_p(\mu_p)$, $V_p \subset E_t$ and $\mathcal{K}(V_p)$ be as in Definition 5. Then for every point o' in a neighborhood of o there are $q(o') \in V_p$ and $\lambda(o')$ (the correspondences $o' \rightarrow q(o')$ and $o' \rightarrow \lambda(o')$ being smooth) such that $o' = \beta_{q(o')}(\lambda_{o'})$. The map $o' \rightarrow (q(o'), \lambda(q(o'))) \in \mathcal{K}(V_p) \subset K(E_t, 2t_*)$ is the desired map. Following the identifications in Definition 5 (which define $K(E_t, 2t_*)$), it is simple to see that the map we defined is indeed independent of the choice of V_p .

With this identification in mind we consider now the set

$$\Omega_{t+2t_*} := (O_{[0,2t_*)}(B_t^2) \cap \Sigma) \cup \overline{\Omega_t},$$

as a set inside $K(E_t, 2t_*)^\circ$, where $\Omega_t = \text{Int}(E_t) \cap \Sigma$. We claim that Ω_{t+2t_*} is a Cauchy surface of the subset F of $K(E_t, 2t_*)^\circ$

$$F = F_1 \cup F_2 \cup F_3,$$

where

$$\begin{aligned} F_1 &= E_t^+, \\ F_2 &= O_{[0,2t_*)}((\partial^N E_t^+) \setminus B_t^2), \\ F_3 &= O_{[0,2t_*)}(B_t^2) \cap \mathcal{M}^+ \end{aligned}$$

and where to define F_2 and F_3 as subsets of $K(E_t, 2t_*)^\circ$ we are using the identification constructed before. To see the claim note first that Ω_t , which is a subset of Ω_{t+2t_*} is a Cauchy surface for E_t^+ . Then by noting that every inextensible past directed causal curve in E_t starting at a point in E_t° cannot reach $\partial^N E_t$, conclude that every inextensible past directed causal curve in F starting at a point in $F_2 \cup F_3$ must either first reach $\partial^N E_t^+$ or eventually reach $\Omega_{t+2t_*} \setminus \Omega_t$. The claim follows.

Now as $\Omega_{t+2t_*} \subseteq \Sigma^\circ$ we have $F \subseteq \mathcal{M}^+$. We obtain therefore $K(E_t^+, t_*) \subset (F \cup O_{[0,2t_*)}(B_t^2))$, but $F \subseteq \mathcal{M}$ and $O_{[0,2t_*)}(B_t^2) \subseteq \mathcal{M}$, thus $K(E_t^+, t_*) \subseteq \mathcal{M}$, which proves **C1**.

We now show **C2**. Suppose **C2** is false. Since by **H2** $T(\partial\Sigma, 2d) \cap E_t = \emptyset$, there exists $0 < \bar{t} < t_*$ such that $J^-(K(E_t, \bar{t}))$ intersects $T(\partial\Sigma, 2d)$ and no smaller $0 < \bar{t} < t_*$ has this

property. Let $p \in \partial T(\partial \Sigma, 2d) \cap J^-(K(E_t, \bar{t}))$ and let $\gamma_p(\tau)$ be the future directed null geodesic on $\partial^N J^-(K(E_t, \bar{t}))$ starting at p . Consider $\gamma_p(\tau'_j)$ where $\tau'_j \rightarrow \infty$ is a divergent sequence. Then we know

1. $(J^-(\gamma_p(\tau'_j)) \cap \Sigma) \subset (J^-(K(E_t, \bar{t})) \cap \Sigma) \subset \Sigma_{2d}$,
2. p is a smooth point of $\partial^N J^-(\gamma_p(\tau'_j))$,
3. $\partial^N J^-(\gamma_p(\tau'_j)) \cap \Sigma$ is tangent to $\partial \Sigma_{2d}$ at p .

Thus a standard comparison of mean curvatures implies that the expansion $\bar{\theta}^+(p)$ of $\partial^N J^-(\{\gamma_p(\tau'_j)\}) \cap \Sigma$ at p is less or equal than the expansion $\theta^+(p)$ of $\partial \Sigma_{2d}$ at p , hence negative. By the Rauchadury equation the foliation of null geodesics of $\partial^N J^-(\{\gamma_p(\tau'_j)\})$ must develop a focussing point along $\gamma_p(\tau)$ in a parametric affine parameter less than a fixed value depending on $\theta^+(p)$. This contradicts the fact that $\tau'_j \rightarrow \infty$ and that $\gamma_p(\tau)$ has no focal points between $\tau = 0$ and $\tau = \tau'_j$.

We show now **C3**. We want to prove

$$(11) \quad J^-(K(E_t^+, t_*)) \cap \mathcal{M}^+ = J^-(\mathcal{W}_{t+t_*}) \cap \mathcal{M}^+$$

The inclusion of the right hand side into the left hand side follows directly because $\mathcal{W}_{t+t_*} \subseteq K(E_t^+, t_*)$. We prove now the inclusion of the left hand side into the right hand side. Let $p \in J^-(K(E_t^+, t_*)) \cap \mathcal{M}^+$. Then there is $q \in K(E_t^+, t_*)$ and a future causal curve γ_1 joining p to q . If $q \in E_t^+$ then we are done as $E_t^+ = J^-(\mathcal{W}_t) \cap \mathcal{M}^+$ and therefore there is a future causal curve γ_2 joining q to a point in $\mathcal{W}_t \subset \mathcal{W}_{t+t_*}$. Thus $\gamma_2 \circ \gamma_1$ (the concatenation of γ_1 and γ_2) is a future causal curve joining p to a point in \mathcal{W}_{t+t_*} . Hence p belongs to the right hand side of (11). If instead $q \in K(E_t^+, t_*) \setminus E_t^+$ then $q = O_{[\bar{t}]}(q')$ where $0 < \bar{t} \leq t_*$ and $q' \in \partial^N E_t^+$. Then there is a null geodesic γ_2 inside $\partial^N E_t^+$ starting at q' and eventually becoming a null geodesic of $\partial^N \mathcal{W}_t$. Therefore $O_{[\bar{t}]}(\gamma_2)$ is a future null geodesic starting at q and eventually becoming a null geodesic of $\partial^N \mathcal{W}_{t+t_*}$. Therefore the curve $O_{[\bar{t}]}(\gamma_2) \circ \gamma_1$ is a future causal curve joining p to a point in \mathcal{W}_{t+t_*} . \square

Proof of Theorem 5: Suppose that $\Sigma^T \setminus \Omega_\infty \neq \emptyset$ (recall we are using $\Omega_\infty = \Sigma \cap E_\infty^+ = \Sigma \cap (\cup E_i^+)$). Let $p \in \partial^T(\Sigma^T \setminus \Omega_\infty)$ (where the topological boundary of $\Sigma^T \setminus \Omega_\infty$ is taken as it were a set in Σ^T). Then $\xi(p)$ is time-like and future directed. Let $\{q_j\} \subset \Sigma^T \cap \Omega_\infty$ be a sequence approaching p , namely $q_j \rightarrow p$. Then, there is $\infty > \lambda_0 > 0$ such that the piece of orbit $\beta_{q_j}(\lambda)$, $\lambda \in (0, \lambda_0]$ lies in \mathcal{M}^+ for all j . Note that for every j there is $i(j)$ such that $q_j \in \Omega_{i(j)}$ (and therefore that $q_j \in E_i^+$ for all $i \geq i(j)$). We claim that the piece of orbit above also lies in E_∞^+ . To see that observe that to leave E_∞^+ it must first leave E_i^+ for all $i \geq i(j)$. That means that for every (i, j) , $i \geq i(j)$ there are $\lambda_{j_i} < \lambda_{j_{i+1}} < \lambda_0$ such that $\beta_{q_j}(\lambda)$, $\lambda \in (\lambda_{j_i}, \lambda_{j_{i+1}})$ lies inside $E_{i+1}^+ \setminus E_i^+$ and that $\beta_{q_j}(\lambda_{j_i}) \in \partial^N E_i^+$ and $\beta_{q_j}(\lambda_{j_{i+1}}) \in \partial^N E_{i+1}^+$. Then, because $E_{i+1}^+ = J^-(K(E_i^+, t_*)) \cap \mathcal{M}^+$, the piece of orbit $\beta_{q_j}(\lambda)$, $\lambda \in (\lambda_{j_i}, \lambda_{j_{i+1}})$ must lie

inside $O_{[0,t_*]}(\partial^N E_i^+) \cap \mathcal{M}^+ \subset E_{i+1}^+$. Therefore it must be $\lambda_{j_{i+1}} - \lambda_{j_i} \geq t_*$. It follows from here that, given j , then $\lambda_{j_i} \rightarrow \infty$ as $i \rightarrow \infty$. Thus it must be $\lambda_0 = \infty$ which is a contradiction, and the orbit $\beta_{q_j}(\lambda)$, $\lambda \in (0, \lambda_0]$ lies in E_∞^+ . Finally we observe that because $\beta_{q_j}(\lambda)$, $\lambda \in [0, \lambda_0]$ is a time-like curve starting at q_j and ending at a point in E_∞^+ , and therefore ending in one of the E_i^+ 's, then if q_j is sufficiently close to p the point p will lie in the interior of Ω_∞ which is against the hypothesis. \square

3 Static Killing initial data

3.1 Background and definitions

We start with the notion of *static Killing initial data*.

Definition 7 A static Killing initial data (static KID) set \mathcal{D} is a KID satisfying the staticity equations

$$(12) \quad NdY + 2Y \wedge Z = 0,$$

$$(13) \quad Y \wedge dY = 0,$$

where $Z := dN + K(Y, \cdot)$.

In a static KID, consider the open set $\Sigma^Y := \Sigma^\circ \setminus \{Y = 0\}$. By the Fröbenius theorem, the distribution Y^\perp is integrable. More precisely, each point $p \in \Sigma^Y$ is contained in a unique, maximal, arc-connected, injectively immersed $(m-1)$ -dimensional, orientable submanifold \mathcal{L}_α orthogonal to Y . The collection of $\{\mathcal{L}_\alpha\}$ is a foliation of Σ^Y . The staticity equation (12) and (4) imply

$$(14) \quad \lambda dY + Y \wedge d\lambda = 0.$$

As a consequence of this equation, if $\lambda = 0$ (resp. $\lambda > 0$, $\lambda < 0$) at any point $p \in \mathcal{L}_\alpha$ then $\lambda = 0$ (resp. $\lambda > 0$, $\lambda < 0$) everywhere on \mathcal{L}_α . To see this, consider any path $\gamma(s)$ contained in \mathcal{L}_α . Contracting (14) with Y and $\dot{\gamma}$ we obtain the ODE

$$\frac{d\lambda(s)}{ds} = Q(s)\lambda(s),$$

where $Q(s)$ is smooth and $\lambda(s) := \lambda(\gamma(s))$. The claim follows.

As discussed in the Introduction, the aim of this part of the work is to show that Killing prehorizons of the exterior region are necessarily embedded. Killing prehorizons are immersed null hypersurfaces where the Killing vector is null and tangent (hence also normal). Thus, their intersection with Σ must correspond to those leaves \mathcal{L}_α where λ vanishes identically. Since we are interested only on horizons of the exterior region or, more precisely,

on horizons than can be reached from the exterior, timelike region, we adopt the following definition.

Definition 8 A horizon \mathcal{H}_α is a leaf of the foliation $\{\mathcal{L}_\alpha\}$ of Σ^Y which intersects the topological boundary $\partial^T \Sigma^T$.

Any two points p_1, p_2 on a fixed leaf \mathcal{L}_α admit transverse sections (i.e. smooth connected curves that are transverse to all the leaves they intersect) Γ_1 and Γ_2 to the foliation $\{\mathcal{L}_\alpha\}$ and a smooth diffeomorphism $\phi : \Gamma_1 \rightarrow \Gamma_2$ such that for any leaf $\mathcal{L}_\beta \in \{\mathcal{L}_\alpha\}$ one has $\phi(\mathcal{L}_\beta \cap \Gamma_1) = \mathcal{L}_\beta \cap \Gamma_2$ (this property is the so-called *transverse uniformity of foliations*, see e.g. Theorem 3, p. 49 in [3]). As a consequence, any horizon \mathcal{H}_α is fully contained in $\partial^T \Sigma^T$.

Since λ vanishes on a horizon, $d\lambda$ is necessarily a normal one-form to \mathcal{H}_α . Consequently there exists a scalar function κ_α on \mathcal{H}_α , called the *surface gravity*, satisfying $d\lambda = 2\kappa_\alpha Y$ on \mathcal{H}_α . It is also convenient to introduce a scalar on Σ defined as

$$(15) \quad I_1 := \frac{1}{4} |dY|_g^2 - 2|Z|_g^2.$$

An alternative expression for I_1 on the set $\{N \neq 0\}$ (in particular on $\{\lambda > 0\}$) is obtained from the staticity equation (12), which gives $dY = -\frac{2}{N}(Y \wedge Z)$. Squaring this and inserting into the definition of I_1 yields

$$(16) \quad I_1 = \frac{2}{N^2} (-\lambda |Z|_g^2 - \langle Y, Z \rangle_g^2) \quad \text{on} \quad \{N \neq 0\}.$$

Passing to the Killing development (which exists in a neighbourhood of any point $p \in \mathcal{H}_\alpha$ since $N|_p \neq 0$), it follows from standard properties of Killing horizons that κ_α is constant on each horizon \mathcal{H}_α (see e.g. [25] p. 334 for a derivation in four dimensions which is, in fact, valid in any dimension).

The following lemma relates I_1 to the surface gravity on horizons.

Proposition 6 $I_1 = -2\kappa_\alpha^2$ on \mathcal{H}_α .

Proof: From (16), it suffices to show that $\kappa_\alpha = \frac{2\langle Y, Z \rangle_g}{N}$. Contracting (4) with Y gives $K(Y, Y) = -\frac{1}{2N} Y(|Y|_g^2)$ which inserted in Z (see Definition 7) gives, on \mathcal{H}_α ,

$$(17) \quad \frac{2\langle Y, Z \rangle_g}{N} = \frac{1}{N^2} Y(N^2 - |Y|_g^2) = \frac{1}{N^2} Y(\lambda) = 2\kappa_\alpha.$$

□

Horizons with non-zero surface gravity have properties qualitatively different to horizons with vanishing surface gravity. The following definition is standard.

Definition 9 A horizon \mathcal{H}_α is degenerate if $\kappa_\alpha = 0$ and non-degenerate if $\kappa_\alpha \neq 0$.

Points where the Killing vector vanishes correspond, at the initial data level, to points $p \in \Sigma$ satisfying $N|_p = Y|_p = 0$. Such points are called *fixed points*. The following lemma is well-known in static four-dimensional spacetimes. At the initial data level, it has been proved in four dimensions in [4]. We include a proof for m -dimensional static KIDs in Appendix A.

Lemma 2 $I_1 < 0$ on any fixed point $p \in \partial^T \Sigma^T$.

In this part of the paper we intend to work directly at the initial data level. This has the advantage that no assumption on well posedness of the matter model needs to be made. Nevertheless, we still require the null energy condition to hold. The following definition translates the standard spacetime definition into the initial data setting.

Definition 10 A Killing initial data set \mathcal{D} satisfies the null energy condition if and only if

$$\mathcal{T}(w, w) - 2J(w)|w|_g + \rho|w|_g^2 \geq 0$$

for any vector $w \in T_p \Sigma$ and $p \in \Sigma$.

3.2 The statements of the main results: Theorem 6 and Corollary 1

Our main result in this second part of the paper is the following.

Theorem 6 Let \mathcal{D} be an asymptotically flat static Killing initial data set satisfying the null energy condition. Suppose that $\partial\Sigma$ (if non-empty) does not intersect Σ^T . Then, each degenerate horizon is an embedded manifold and compact.

An immediate Corollary of Theorems 5 and 6 is

Corollary 1 Let \mathcal{D} be an asymptotically flat static Killing initial data set with well posed matter model satisfying the null energy condition. Suppose that $\partial\Sigma$ (if non-empty) is future outer trapped. Then each degenerate horizon is an embedded manifold and compact.

Remark 2 It may be possible to prove, directly from the techniques that we developed here, a version of Theorem 6 also for stationary data and not just static. We will not enter into such problem here however.

3.3 Volume monotonicity along “optic” congruences of geodesics

In this section, we will assume that the datum \mathcal{D} is static (Definition 7).

The Killing development of a static KID is static in the sense that the Killing vector ξ is hypersurface orthogonal (see Lemma 3 in [5]). Static spacetimes necessarily satisfy

$\mathbf{G}(\xi, X) = 0$, where X is any vector field orthogonal to ξ . In terms of the quantities (ρ, J, \mathcal{T}) defined by (1), this implies

$$(18) \quad \mathcal{T}(Y, \cdot) = NJ + \frac{J(Y)}{N}Y - \rho Y, \quad \text{on } \{N \neq 0\} \subset \Sigma.$$

In addition to g , Σ^T can be endowed with two further metrics: the so-called *quotient metric*

$$(19) \quad h := g + \frac{1}{\lambda}Y \otimes Y$$

and the *optic metric*

$$(20) \quad \bar{h} := \frac{1}{\lambda}h.$$

Consider the spacetime $(\Sigma^T \times \mathbb{R}, g_S)$ with metric

$$(21) \quad g_S = -V^2 dt'^2 + h$$

where $V := +\sqrt{\lambda}$. Equation (14) implies that $\lambda^{-1}Y$ is closed on open sets where λ does not vanish, in particular on Σ^T . Consequently there exists, locally, a function ζ such that $Y = -\lambda d\zeta$. The coordinate transformation $t = t' - \zeta$ brings the metric g_D (see (8)) into g_S . This shows that the spacetimes $(\Sigma^T \times \mathbb{R}, g_D)$ and $(\Sigma^T \times \mathbb{R}, g_S)$ are locally isometric. They are also globally isometric if $\lambda^{-1}Y$ is exact on Σ^T .

Since the data on $\{t' = 0\}$ in the metric (21) is a totally geodesic static KID, it satisfies the constraint equations (2)-(3) and the KID equations (4)-(5) with the substitutions $g \rightarrow h$, $N \rightarrow V$, $Y \rightarrow 0$ and $K \rightarrow 0$. With the definitions $\hat{\rho} := V^{-2}\mathbf{G}(\xi, \xi)$ and $\hat{\mathcal{T}}(v, w) := \mathbf{G}(v, w)$, with v, w tangent to $\{t' = 0\}$, these equations read

$$(22) \quad \text{Hess}_h V = V \left(\text{Ric}_h - \hat{\mathcal{T}} + \frac{1}{m-1} (\text{tr}_h \hat{\mathcal{T}} - \hat{\rho}) h \right),$$

$$(23) \quad \Delta V = V \left(\frac{\text{tr}_h \hat{\mathcal{T}}}{m-1} + \frac{m-2}{m-1} \hat{\rho} \right),$$

where Δ is the Laplacian of h and Ric_h its Ricci tensor. Using $\xi = Nn + Y$ and (1), it is straightforward to relate $\hat{\rho}$, $\hat{\mathcal{T}}$ to the datum \mathcal{D} , as follows

$$(24) \quad \hat{\rho} = \rho - \frac{1}{N}J(Y), \quad \hat{\mathcal{T}} = \mathcal{T} + \frac{1}{\lambda} \left(\frac{J(Y)}{N} - \rho \right) Y \otimes Y.$$

The following Proposition characterizes the null energy condition of the initial data set in terms of the geometry associated to h .

Proposition 7 \mathcal{D} satisfies the null energy condition if and only if

$$\hat{\mathcal{T}}(\hat{w}, \hat{w}) + \hat{\rho}|\hat{w}|_h^2 \geq 0$$

for any vector $\hat{w} \in T_p \Sigma^T$ and $\forall p \in \Sigma^T$.

Proof: This Proposition can be proved easily by passing to the Killing development of Σ^T . For a direct proof on the initial data set, consider any vector \hat{w} and define $w := \hat{w} + \frac{A}{V}Y$, where $A := |\hat{w}|_h + \frac{\langle Y, \hat{w} \rangle_g}{V}$. The g -norm of w is calculated to be $|w|_g^2 = \frac{N^2 A^2}{\lambda}$. A straightforward computation which uses (24) and (18) gives

$$\mathcal{T}(w, w) - 2J(w)|w|_g + \rho|w|_g^2 = \hat{\mathcal{T}}(\hat{w}, \hat{w}) + \hat{\rho}|\hat{w}|_h^2.$$

Since transformation $\hat{w} \rightarrow w$ is invertible (with inverse $\hat{w} = w - \frac{|w|_g}{N}Y$), the Proposition follows from Definition 10. \square

Expression (22) determines the Ricci tensor of h in terms of V and its derivatives. A similar expression can be obtained for the Ricci tensor of \bar{h} , denoted by $Ric_{\bar{h}}$. We write ∇ for the covariant derivative of h and $\bar{\nabla}$ for the covariant derivative of \bar{h} .

Proposition 8 The Ricci tensor of \bar{h} takes the following form

$$(25) \quad Ric_{\bar{h}} = (m-1)\frac{1}{V}Hess_h V - (m-1)\frac{|\nabla V|_h^2}{V^2}h + \hat{\rho}h + \hat{\mathcal{T}}.$$

Proof: The general expression for the change of Ricci tensor under a conformal rescaling $\bar{h} = e^{2f}h$ is

$$Ric_{\bar{h}} = Ric_h + (2-m)(Hess_h f - df \otimes df) - (\Delta f + (m-2)|\nabla f|_h^2)h.$$

Putting $f = -\ln(V)$ and inserting (22) and (23), the Proposition follows. \square

The following proposition is well-known [26] and explains the reason of calling \bar{h} the *optic metric*.

Proposition 9

1. Let $\gamma(t)$, $t \in [t_0, t_1]$ be a geodesic segment in (Σ^T, \bar{h}) parametrized by \bar{h} -arc-length. Select $c \neq 0$, define

$$\tau(t) = \tau_0 + \int_{t_0}^t c^{-1} V^2(\gamma(t)) dt$$

and denote by $t(\tau)$ its inverse (which obviously exists). Then the curve $(\gamma(t(\tau)), t(\tau))$, $\tau \in [\tau_0, \tau(t_1)]$ is an affinely parametrized null geodesic segment in $(\Sigma^T \times \mathbb{R}, g_S)$ and its tangent vector v satisfies $g_S(v, \xi) = -c$.

2. Conversely, let $(\gamma(\tau), t(\tau))$, $\tau \in [\tau_0, \tau_1]$ be an affinely parametrized null geodesic segment in $(\Sigma \times \mathbb{R}, g_S)$ with tangent vector v . Define $c = -g_S(v, \xi)$ (which is obviously constant along the geodesic) and define $\tau(t)$ as the inverse of

$$t(\tau) := t_0 + \int_{\tau_0}^{\tau} \frac{c}{V^2(\gamma(\tau))} d\tau.$$

Then, the curve $\gamma(\tau(t))$, $t \in [t_0, t(\tau_1)]$ is a geodesic segment in (Σ^T, \bar{h}) parametrized by \bar{h} -arc-length.

Consider a smooth, oriented hypersurface S embedded in Σ^T and let ν and $\bar{\nu}$ be positively oriented normal vectors, of unit length respectively in h and \bar{h} (they are obviously related by $\bar{\nu} = V\nu$). Let h_S (resp. \bar{h}_S) denote the induced metric on S inherited from h (resp. \bar{h}). The following fact is well-known and straightforward.

Proposition 10 *With the notation before, the second fundamental form χ of S with respect to ν in the metric h and the second fundamental form $\bar{\chi}$ of S with respect to $\bar{\nu}$ in the metric \bar{h} are related by*

$$\bar{\chi} = \frac{\chi}{V} - \frac{\nu(V)}{V^2} h_S.$$

Squaring in their respective metrics and taking traces the following expressions follow,

$$(26) \quad \begin{aligned} \bar{\theta} &= V\theta - (m-1)\nu(V), \\ |\bar{\chi}|_{\bar{h}_S}^2 &= V^2 |\chi|_{h_S}^2 + \frac{V^2}{m-1} \left(\theta - \frac{(m-1)}{V} \nu(V) \right)^2, \end{aligned}$$

where $\theta := \text{tr}_{h_S} \chi$, $\bar{\theta} := \text{tr}_{\bar{h}_S} \bar{\chi}$ and Π is the trace-free part of χ (in the metric h_S). The expression above for $\bar{\theta}$ and the Ricci tensor of \bar{h} give rise to the following monotonicity formula.

Proposition 11 *Let \mathcal{F} be a congruence of geodesics in (Σ^T, \bar{h}) parametrized by arc-length. Assume that the tangent vector $\bar{\nu}$ to this congruence is orthogonal to a collection of smooth hypersurfaces $\{S_t\}$. Then, the trace θ of the second fundamental form χ of S_t with respect to ν satisfies*

$$(27) \quad \bar{\nu} \left(\frac{\theta}{V} \right) + |\chi|_{h_S}^2 + \frac{1}{m-1} \theta^2 + \hat{\rho} + \hat{\mathcal{T}}(\nu, \nu) = 0.$$

In particular, if the null energy condition is satisfied in (Σ^T, h) then

$$\bar{\nu} \left(\frac{\theta}{V} \right) \leq -\frac{1}{m-1} \theta^2 \leq 0.$$

Proof: The focusing equation for geodesics (see e.g. [7]) is

$$(28) \quad \bar{v}(\bar{\theta}) + |\bar{\chi}|_{\bar{h}_g}^2 + Ric_{\bar{h}}(\bar{v}, \bar{v}) = 0.$$

The term $Ric_{\bar{h}}(\bar{v}, \bar{v})$ can be directly evaluated from (25):

$$(29) \quad Ric_{\bar{h}}(\bar{v}, \bar{v}) = (m-1)V \text{Hess}_h V(\nu, \nu) - (m-1)|\nabla V|_h^2 + V^2(\hat{\rho} + \hat{\mathcal{T}}(\nu, \nu)).$$

In order to evaluate the term $\bar{v}(\bar{\theta})$ in (28), the h -acceleration $\nabla_\nu \nu$ is needed. Since \bar{v} is geodesic and affinely parametrized we have $\bar{\nabla}_{\bar{v}} \bar{v} = 0$, which becomes, after applying the transformation law for metric connections under conformal rescalings,

$$(30) \quad \nabla_\nu \nu = \frac{1}{V} \nu(V) \nu - \frac{1}{V} \nabla(V).$$

We then have, from (26),

$$(31) \quad \begin{aligned} \bar{v}(\bar{\theta}) &= \bar{v}(V)\theta + V \bar{v}(\theta) - (m-1)V [\langle \nabla_\nu \nu, \nabla V \rangle_h + (\text{Hess}_h V)(\nu, \nu)] \\ &= \bar{v}(V)\theta + V \bar{v}(\theta) + (m-1) \left(|\nabla V|_h^2 - \nu(V)^2 - V(\text{Hess}_h V)(\nu, \nu) \right). \end{aligned}$$

Inserting (29) and (31) into (28), the terms in the Hessian of V cancel out. A simple rearrangement gives (27). The last claim follows from Proposition 7. \square

3.4 On the volume of horizons of asymptotically flat static KIDs

Recall that Σ^∞ is the AF end of \mathcal{D} . The decay (7) implies that (Σ^∞, h) is also asymptotically flat. Let S_r and $\Sigma^I(r)$ be defined as in Section 2.1 and define $\Sigma_I^T := \Sigma^I(r) \cap \Sigma^T$. We start by showing that Σ_I^T is complete in the metric \bar{h} .

Lemma 3 *Assume that Σ^T does not intersect $\partial\Sigma$. Then, the Riemannian manifold (Σ_I^T, \bar{h}) is complete and has S_r as its only boundary.*

Proof: First we make a couple of comments on the structure of the metric \bar{h} around (I) a point on a horizon and (II) a fixed point.

(I) Consider a point p lying on a horizon \mathcal{H}_α and choose a foliated chart $(V_p, \{x^A, z\})$ near p adapted to the foliation $\{\mathcal{L}_\alpha\}$. This means that, in these coordinates, $V_p = \Omega \times (-\delta, \delta)$, where Ω is a domain on \mathbb{R}^{m-1} and $\delta > 0$. The coordinate z takes values in $(-\delta, \delta)$ and $\{x^A\}$ ($A, B = 1, \dots, m-1$) takes values in Ω . The intersection of any leaf \mathcal{L}_α with V_p is a collection (possibly empty) of sets of the form $\Omega \times \{z_i\}$ (called plaques) where $\{z_i\}$ is a countable set. Since Y is g -orthogonal to the plaques, we can choose, without loss of generality, the coordinate chart so that the metric g takes the form $g = F^2(z, x^C) dz^2 + \hat{g}_{AB}(z, x^C) dx^A dx^B$, where $F > 0$ and \hat{g}_{AB} is positive definite. Furthermore, we can assume that $p = \{0\}$. Let P be the smooth positive function on V_p such that $Y = NP\nu$ where ν is

g -unit and orthogonal to the plaques. This implies $\lambda = N^2(1 - P^2)$. Inserting all this into (14) yields, after a straightforward calculation,

$$(1 - P^2) \left(-\frac{\partial_A F}{F} + \frac{\partial_A N}{N} \right) - \frac{1 + P^2}{P} \partial_A P = 0, \quad \text{for all } z \in (-\delta, \delta)$$

which, upon integration, implies the existence of a smooth function U on V_p , constant on every plaque (i.e. $U(z)$), such that

$$\frac{N}{F} \frac{1 - P^2}{P} = U.$$

Using this into the definition of \bar{h} (20) gives

$$(32) \quad \bar{h} = \frac{dz^2}{P^2 U(z)^2} + \frac{1}{N F P U(z)} \hat{g}_{AB} dx^A dx^B.$$

Note that from $\lambda|_p = 0$ we have $U(z = 0) = 0$. Also, as U is differentiable, we have (on Σ^T) $0 < U \leq c|z|$ (where $c > 0$ is a constant) near p .

(II) Let now p be a fixed point. Then we know from (15) and Lemma 2 that $Z|_p \neq 0$. This in turn implies $dN|_p \neq 0$ (see Definition 7). Thus, there exists a neighbourhood V_p of p where N can be taken as a coordinate. Without loss of generality, we can choose a coordinate system in V_p so that $g = \tilde{F}^2(N, x^C) dN^2 + \tilde{g}_{AB}(N, x^C) dx^A dx^B$ where $\tilde{F} > 0$ and \tilde{g}_{AB} is positive definite. By the definition of \bar{h} , we have

$$(33) \quad \bar{h} \geq \frac{1}{\lambda} g \geq \frac{1}{N^2} g.$$

Note that $N|_p = 0$ and that, once more, N is a coordinate in a differentiable coordinate system.

We are ready to prove completeness of (Σ_I^T, \bar{h}) . Assume by contradiction that \bar{h} is not complete. Let γ be an incomplete \bar{h} -geodesic not ending at S_r . Then γ , as a curve over Σ accumulates (although not necessarily converging to) a point p on a horizon or a fixed point p . From the structure of the metric \bar{h} found around such points in (32) and (33), respectively, one readily deduces that the \bar{h} -length of γ must be infinite which is a contradiction. \square

On Σ_I^T define $B(t, S_r)$, $t > 0$, as the \bar{h} -ball of center S_r and radius t ,

$$B(t, S_r) = \{p \in \Sigma_I^T, \text{dist}_{\bar{h}}(p, S_r) < t\}.$$

The boundary component

$$\partial B(t, S_r) := \partial^T B(t, S_r) \setminus S_r$$

is the set of points lying at \bar{h} -distance t to S_r . Outside the cut locus C this set of points is a smooth hypersurface. We want to consider the $(m-1)$ -Hausdorff measure of $\partial B(t, S_r)$ in the metric h , which we denote by $|\partial B(t, S_r)|_h$. The following lemma gives an upper bound for

$|\partial B(t, S_r)|_h$.

Lemma 4 *Let S_r be the coordinate sphere of radius r in Σ^∞ and assume that the h -mean curvature with respect to the ingoing unit vector is negative everywhere. Assume that Σ^T does not intersect $\partial\Sigma$ and let $|S_r|_h$ be the $(m-1)$ -volume of S_r in the metric h . Then, $|\partial B(t, S_r)|_h \leq |S_r|_h$ for all $t > 0$.*

Proof: On Σ_I^T consider the congruence \mathcal{F} of geodesics minimizing the \bar{h} -distance to S_r . An immediate consequence of Lemma 3 is that each geodesic in \mathcal{F} has an end-point in S_r . For any $p \in \Sigma_I^T$ outside the cut locus C of the distance function (which has zero measure [19]) the function $t(p) = \text{dist}_{\bar{h}}(p, S_r)$ is smooth and in there the level sets of t are smooth hypersurfaces. In other words, if $p \in \Sigma_I^T \setminus C$ then, around p , $\partial B(t(p), S_r)$ is a smooth hypersurface. Let p be such a point and let $\gamma_p(t)$ be the length minimizing segment that starts at S_r and ends at p . We define the function θ on $\Sigma^T \setminus C$ as the h -mean curvature of $\partial B(t(p), S_r)$ at p in the direction of $\gamma'_p(t(p))$. Note that the mean curvature is with respect to h and not \bar{h} , but that the congruence \mathcal{F} is with respect to \bar{h} and not h .

Now, from Proposition 11 we have the monotonicity

$$\bar{v}\left(\frac{\theta}{V}\right) \leq -\frac{\theta^2}{m-1} \leq 0.$$

Since $\theta|_{S_r} < 0$ we conclude that $\theta < 0$ on $\Sigma^T \setminus C$. Denoting by $\eta_h(p)$ the volume-form of $\partial B(t(p), S_r)$ at $p \in \Sigma^T \setminus C$, the first variation $(m-1)$ -volume gives

$$\bar{v}(\eta_h) = Vv(\eta_h) = V\theta < 0.$$

This proves $|\partial B(t, S_r)|_h \leq |S_r|_h$. □

We analyze now the interplay between the $(m-1)$ -volume of horizons in the static KID and the $(m-1)$ -volume of the \bar{h} -geodesic spheres $\partial B(t, S_r)$.

Let \mathcal{H}_α be a horizon and let v be one of the two possible normal vector fields to \mathcal{H}_α . For every point $q \in \mathcal{H}_\alpha$ consider the g -geodesic $\gamma_q(s)$ starting at q with velocity $v(q)$ and parametrized with arc-length. Let $\Omega \subset \mathcal{H}_\alpha$ be open and connected with smooth and compact boundary in \mathcal{H}_α .

Definition 11 *Let \mathcal{H}_α be a horizon. We say that \mathcal{H}_α is isolated on Ω in the direction of v if for some \bar{s} small, the set (tubular neighborhood of Ω)*

$$U_v(\Omega, \bar{s}) = \{\gamma_q(s), q \in \Omega, 0 < s < \bar{s}\},$$

is contained in Σ^T and does not intersect any horizon. A horizon \mathcal{H}_α is isolated if there exists an exhaustion $\{\Omega_i\}$ of \mathcal{H}_α such that \mathcal{H}_α is isolated on Ω_i in both normal directions.

Since $d\lambda \neq 0$ everywhere on a non-degenerate horizon, it follows that non-degenerate horizons are necessarily isolated.

Proposition 12 *Let \mathcal{H}_α be an isolated horizon in the direction of v over Ω . Then*

$$\liminf_{t \rightarrow \infty} |U_v(\Omega, \bar{s}) \cap \partial B(t, S_r)|_h \geq |\Omega|_g.$$

where $|\Omega|_g$ is the g -($m-1$)-volume of Ω .

Proof: We need several definitions first.

1. At every point $p \in U_v(\Omega, \bar{s})$, let v be the tangent of the geodesic $\gamma_q(s)$ passing through p . Choose $(m-1)$ vector fields $\{e_1, \dots, e_{m-1}\}$ on $U_v(\Omega, \bar{s})$ such that $\{e_1, \dots, e_{m-1}, v\}$ is an oriented g -orthonormal basis. Let $\{\omega^1, \dots, \omega^m\}$ be the corresponding dual basis. Define then the $(m-1)$ -form

$$\omega = \omega^1 \wedge \dots \wedge \omega^{m-1}.$$

2. For every $0 < \tilde{s} < \bar{s}$ define the surface

$$S(\Omega, \tilde{s}) = \{\gamma_q(\tilde{s}), q \in \Omega\},$$

and its one-sided tubular neighbourhoods

$$U^+(\Omega, \tilde{s}) = \{\gamma_q(\tilde{s}), q \in \Omega, \tilde{s} \leq s < \bar{s}\}, \quad U^-(\Omega, \tilde{s}) = \{\gamma_q(\tilde{s}), q \in \Omega, 0 < s \leq \tilde{s}\}.$$

Now, for every \tilde{s} there is $t_0(\tilde{s})$ such that if $t > t_0(\tilde{s})$ then $U^+(\Omega, \tilde{s}) \subset B(t, S_r)$, namely $U^+(\Omega, \tilde{s})$ lies in the interior of the \bar{h} -metric ball $B(t, S_r)$. For such t define $B(t, \tilde{s})$ as the connected component of $U_v(\Omega, \tilde{s}) \cap B(t, S_r)$, containing $U^+(\Omega, \tilde{s})$. Then $\partial^T(B(t, \tilde{s}) \setminus U^+(\Omega, \tilde{s}))$ consists of:

1. $S(\Omega, \tilde{s})$,
2. an interior component that we will denote $S_t(\Omega)$ which is in fact equal to a component of $U^-(\Omega, \tilde{s}) \cap \partial B(t, \tilde{s})$, and,
3. a domain inside the $(m-1)$ -surface $\{\gamma_g(s), q \in \partial\Omega, 0 < s < \tilde{s}\}$.

Since the metric h is related to g by (19) their volume forms are related by

$$\eta_h = (1 + \frac{|Y|_g^2}{\lambda}) \eta_g.$$

Consequently

$$|S_t(\Omega)|_h \geq |S_t(\Omega)|_g.$$

On the other hand we have

$$|S_t(\Omega)|_g \geq \int_{S_t(\Omega)} \omega = |S(\Omega, \tilde{s})|_g + \int_{(B(t, \tilde{s}) \setminus U^+(\Omega, \tilde{s}))} d\omega.$$

Integration by parts is justified (for almost all t) because the distance function is Lipschitz and therefore of bounded variation [16] (indeed it is semiconcave and therefore a $H^{2,1}$ function [19]). But now, as $\tilde{s} \rightarrow 0$ and $t > t_0(\tilde{s}) \rightarrow \infty$, the first term on the right hand side approaches $|\Omega|_g$ and the second converges to zero. Since obviously $|U_\nu(\Omega, \tilde{s}) \cap \partial B(t, S_r)|_h \geq |S_t(\Omega)|_h$ we conclude

$$\liminf_{t \rightarrow \infty} |U_\nu(\Omega, \tilde{s}) \cap \partial B(t, S_r)|_h \geq \liminf_{t \rightarrow \infty} |S_t(\Omega)|_h \geq |\Omega|_g.$$

□

Proposition 13 *Assume that $\Sigma^T \cap \partial \Sigma = \emptyset$. Let $\{H_\alpha\}_{\alpha \in \mathcal{J}}$ be any collection of horizons in an asymptotically flat static KID \mathcal{D} . Let $H = \bigcup_{\alpha \in \mathcal{J}} H_\alpha$ be its union. Then $\overline{H} \setminus H$ is either empty or consists only of fixed points.*

Proof: The proof is by contradiction. We will assume that there exists $p \in \overline{H} \setminus H$ which is not a fixed point and we will show that $\lim_{t \rightarrow \infty} |\partial B(t, S_r)|_h = +\infty$, which contradicts the upper bound found in Lemma 4.

Let p be such a point. Since p is non-fixed ($Y|_p \neq 0$), there exists a unique leaf \mathcal{L}_β containing p . Since $H \subset \partial^T \Sigma^T$ (recall that a horizon is fully contained in $\partial^T \Sigma^T$) and the latter is topologically closed, it follows that $p \in \partial^T \Sigma^T$, so in fact \mathcal{L}_β is a horizon \mathcal{H}_β . By hypothesis, this horizon is not in the original collection $\{H_\alpha\}_{\alpha \in \mathcal{J}}$. Consider a foliated chart V_p of p in Σ^Y as in the proof of Lemma 3. Without loss of generality, we can assume that the foliated chart is centered at \mathcal{H}_β , i.e. that the plaque $\Omega \times \{0\} \subset \mathcal{H}_\beta$. Also without loss of generality we assume that $\overline{\Omega}$ is compact with smooth boundary. By definition of horizon, there exists a sequence of points $p_i \rightarrow p$ with $p_i \in \Sigma^T$ (in particular $\lambda(p_i) > 0$). Moreover, since $p \in \overline{H} \setminus H$, there must exist a sequence of plaques in H converging to $\Omega \times \{0\}$. These two facts together imply the existence of two sequences $a_i \rightarrow 0$, $b_i \rightarrow 0$, $-\delta < a_i < b_i < \delta$ such that

1. $\Omega \times (a_i, b_i) \subset \Sigma^T$,
2. $\Omega \times \{a_i\} \in \partial^T \Sigma^T$,
3. $\Omega \times \{b_i\} \in \partial^T \Sigma^T$.

Let \mathcal{H}_i^+ be the horizon containing the plaque $\Omega \times \{b_i\}$, and \mathcal{H}_i^- be the horizon containing the plaque $\Omega \times \{a_i\}$. Both horizons are isolated on Ω (the first one in the direction of decreasing z and the second one towards increasing z). For each t sufficiently large, let $C(t)$ be the maximum of $i \in \mathbb{N}$ such that, $\forall j \leq i$, $\Omega \times (a_j, b_j) \cap B(t, S_r)$ is non-empty and has more than




Figure 2: Foliated neighbourhood of Ω . The grey regions lie in Σ^T . Schematic plots of the balls $B(t, S_r)$ are shown for two values of t , namely t_1 and t_2 satisfying $t_1 < t_2$. As t increases the balls increase and approach the boundaries of Σ^T . Two values of the sequences $\{a_i\}$ and $\{b_i\}$ defined in the main text are also shown. In the case of the figure $C(t_1) = 0$ and $C(t_2) = 1$.

one connected component. By Proposition 12, each one of the pieces $\partial B(t, S_r) \cap (\Omega \times (a_i, b_i))$, $i \leq C(t)$ contributes to the total $(m-1)$ -volume $|\partial B(t, S_r)|_h$ essentially with an amount of at least $2|\Omega|_g$. More precisely, for fixed $\epsilon > 0$, there exists $t_0(\epsilon)$ such that for $t > t_0(\epsilon)$

$$|\partial B(t, S_r)|_h \geq 2C(t) (|\Omega|_g - \epsilon)$$

Since $C(t) \rightarrow \infty$ when $t \rightarrow \infty$, we obtain $\lim_{t \rightarrow \infty} |\partial B(t, S_r)|_h = +\infty$ and hence a contradiction to Lemma 4. \square

We can now prove Theorem 6.

Proof of Theorem 6: We first show that any degenerate horizon \mathcal{H}_α is compact. I_1 vanishes on \mathcal{H}_α and hence also on its closure. Assume that there is a point $p \in \overline{\mathcal{H}_\alpha} \setminus \mathcal{H}_\alpha$. p must be a fixed point by Proposition 13. However, since $\mathcal{H}_\alpha \in \partial^T \Sigma^T$, it follows $p \in \partial^T \Sigma^T$. Lemma 2 gives $I_1 < 0$, which gives a contradiction. Hence \mathcal{H}_α is topologically closed. Closed leaves in foliations are necessarily embedded (see e.g. Theorem 5, p. 51 in [3]). Moreover, since \mathcal{H}_α is contained in the compact set $\Sigma^I(r)$, \mathcal{H}_α is also compact. \square

The previous results prove not only that a degenerate horizon cannot approach itself indefinitely, but also that two or more such horizons cannot wrap on themselves indefinitely. More precisely, we have the following Corollary.

Corollary 1 *Assume that $\Sigma^T \cap \partial \Sigma = \emptyset$. Then all horizons are isolated. Moreover $|\bigcup_\alpha \mathcal{H}_\alpha|_g < |S_r|_h$.*

Proof: Assume that \mathcal{H}_α is a non-isolated horizon, hence necessarily degenerate. Since \mathcal{H}_α is not isolated, there exists an open set $\Omega \subset \mathcal{H}_\alpha$ with compact, smooth boundary in \mathcal{H}_α such that for all $\bar{s} > 0$, the tubular neighbourhood

$$U(\Omega, \bar{s}) = \{\gamma_q(s), q \in \Omega, -\bar{s} < s < \bar{s}\},$$

intersects another horizon \mathcal{H}_β . Selecting a sequence $\bar{s}_i \rightarrow 0$, we have a collection of horizons \mathcal{H}_{β_i} which approach \mathcal{H}_α . By Theorem 6 \mathcal{H}_α is embedded. It follows that at least one of the $\mathcal{H}_{\beta_i} \neq \mathcal{H}_\alpha$. Consider the collection \mathcal{A} of all $\{\mathcal{H}_{\beta_i}\}$ different from \mathcal{H}_α . It follows that the set $H := \cup_{\mathcal{A}} \mathcal{H}_{\beta_i}$ is not closed, as its closure contains Ω . This contradicts Proposition 13. Thus \mathcal{H}_α is isolated. The last statement is a direct consequence of Proposition 12 and Lemma 4. \square

4 Uniqueness of static, vacuum, asymptotically flat initial data sets with outer trapped boundary

The results of the previous sections allow us to prove a uniqueness theorem for asymptotically flat static KID with an outer trapped boundary.

The most powerful method of proving uniqueness of static black holes is the so-called *doubling method* of Bunting and Masood-ul-Alam [2]. The framework where this method applies involves asymptotically flat KID such that the exterior region Σ^T where the Killing vector is timelike has a topological boundary which is a compact, embedded C^0 manifold without boundary (see [11] for details in the vacuum case). More generally, the method applies also to settings where the Cauchy boundary of Σ^T is a compact, embedded C^0 manifold without boundary (the Cauchy boundary is defined as the set of points lying in the Cauchy completion of the set but not in the set itself). The need of using Cauchy completions comes from the fact that a point $p \in \partial^T \Sigma^T$ may be accessible from both sides from within Σ^T . In this setting $\partial^T \Sigma^T$ may not be a C^0 manifold but the Cauchy boundary, denoted by $\partial^C \Sigma^T$ may still be a topological manifold. We will see an example of this behavior later.

A possible strategy for proving a uniqueness theorem for static KID with an outer trapped boundary is to reduce the problem to a black hole uniqueness theorem. This suggests the following definition (c.f. [4, 6]): *an asymptotically flat KID \mathcal{D} (possibly with boundary) is a black hole static initial data set if the Cauchy boundary $\partial^C \text{Int}(\Sigma^T)$ of Σ^T is a topological manifold without boundary and compact.*

In agreement with the discussion above, we will also say that a *matter model satisfies the static black hole uniqueness theorem* if there exists a class of asymptotically flat static spacetimes $\{(M_a, g_a)\}$ depending on a finite (and usually small) number of parameters determined from the asymptotic form of the metric and matter fields and such that any *black hole static Killing initial data set* \mathcal{D} has the property that (Σ^T, g, K) can be isometrically embedded in some $\{(M_a, g_a)\}$ within this class (where isometrically embedded is in the sense of spacetime

initial data sets).

As example of matter models satisfying the static black hole theorem we have vacuum or electrovacuum in four spacetime dimensions (see [11], [14], [8] and references therein).

An interesting consequence of the results in the previous chapters is that the static black hole uniqueness theorem can be extended to static, asymptotically flat KID with outer trapped boundary. More precisely

Theorem 7 *Let \mathcal{D} be a static, m -dimensional ($m \geq 3$) asymptotically flat Killing initial data satisfying the following assumptions:*

- A1.** *The matter model is well posed and satisfies the null energy condition.*
- A2.** *The matter model satisfies the static black hole uniqueness theorem.*
- A3.** *Σ has outer trapped boundary, i.e. $\partial\Sigma$ is compact and $\theta^+(\partial\Sigma) < 0$.*

Then (Σ^T, g, K) can be isometrically embedded in some (\mathcal{M}_a, g_a) within the black hole uniqueness class.

Remark 3 *In the particular case of vacuum, this result takes the form of Theorem 3 given in the Introduction.*

We first recall a well-known property of fixed points of Killing vectors in spacetimes of arbitrary dimension.

Lemma 5 *Let (\mathcal{M}, g) be a spacetime with a Killing vector ξ . Let p be a fixed point of ξ , i.e. $\xi(p) = 0$ and let $F := \frac{1}{2}d\xi$ (where $\xi = g(\xi, \cdot)$) be the so-called Killing form of ξ . Define $W_p := \{v \in T_p\mathcal{M}; F|_p(v, \cdot) = 0\}$. Then p lies in smooth, totally geodesic embedded submanifold S_p of dimension $k = \dim(W_p)$ (p is an isolated fixed point if $k = 0$). Moreover, S_p is spacelike, null or timelike depending on whether W_p is spacelike, null or timelike.*

Proof of Theorem 7: From Theorem 5 it follows that Σ^T does not intersect $\partial\Sigma$. By Theorem 6 each degenerate horizon of Σ^T is compact and embedded. The same is true for non-degenerate horizons. Consider any point p in a horizon. Then, near p the Cauchy boundary $\partial^C\Sigma^T$ either coincides with $\partial^T\Sigma^T$ (if Σ^T only lies to one side of $\partial^T\Sigma^T$ at p) or with two copies of $\partial^T\Sigma$ (if p can be accessed from both sides within Σ^T). In either case, the Cauchy completion is a smooth manifold near p .

It only remains to analyze the fixed points $p \in \partial\Sigma^T$. We know from the proof of Lemma 2 in Appendix A that $dN|_p \neq 0$ and that $dY|_p = \frac{2b}{Q}(dN|_p \wedge X)$ for some $b, Q > 0$ and $X \in T_p^*\Sigma$ which is both unit and orthogonal to $dN|_p$. Moreover, $b^2 < Q^2$ from (35) in the proof of Lemma 2. Let us now view p as a point in \mathcal{M} . It is clear that p is a fixed point for the Killing vector ξ . The Killing form F at p is easily evaluated to be (c.f. Definition 3 in [5])

$$F|_p = (dN \wedge n)|_p + \frac{b}{Q}(dN|_p \wedge X),$$

where \mathfrak{n} is the future directed unit normal one-form to Σ in \mathcal{M} . Since the one-form $(\mathfrak{n}|_p + bQ^{-1}X)$ is timelike (from $Q^2 > b^2$), it follows that dN_p and $\mathfrak{n}|_p + bQ^{-1}X$ span a timelike two-plane. From the definition of W_p in Lemma 5 we conclude that W_p is $(m-1)$ -dimensional and spacelike. Thus p lies on a smooth codimension-two, totally geodesic spacelike surface of \mathcal{M} . In these circumstances, the same construction performed by Rácz-Wald [23] in dimension four in order to find a canonical coordinate system near p applies to arbitrary dimension. This gives a coordinate system $\{u, v, x^a\}$ ($a, b = 2, \dots, m$) in an open connected neighbourhood U_p of p with the following properties:

- The metric takes the form

$$g = 2Gdudv + 2vH_adx^adu + g_{ab}dx^adx^b$$

with G, H_a, g_{ab} smooth functions of (uv, x^a) , $G > 0$ and g_{ab} positive definite.

- The surface $S_p \cap U_p$ takes the local form $\{u = 0, v = 0\}$.
- The Killing vector ξ reads $\xi = u\partial_u - v\partial_v$.
- ∂_v is future directed everywhere.

Since ∂_v is null and non-zero, the spacelike hypersurface $\Sigma \cap U_p$ can be written as a graph $\{v = \phi(u, x^a)\}$ (in particular, $\{u, x^a\}$ defines a local coordinate system on $\Sigma \cap U_p$). Since Σ is spacelike ϕ satisfies $\partial_u \phi > 0$ everywhere. Let $u_0(x^a)$ be the solution of $\phi(u, x^a) = 0$ (which exists because ϕ vanishes on p).

Now, since $\lambda = 2\hat{G}u\phi$ where $\hat{G} = G(u\phi, x^a)$ it follows that either

- (i) $\Sigma^T \cap U_p = \{u > 0\} \cap \{u > u_0\}$, or
- (ii) $\Sigma^T \cap U_p = \{u < 0\} \cap \{u < u_0\}$, or
- (iii) $\Sigma^T \cap U_p = (\{u > 0\} \cap \{u > u_0\}) \cup (\{u < 0\} \cap \{u < u_0\})$.

The corresponding Cauchy boundaries are:

For (i): $\{u = \max(0, u_0(x^a)), x^a\}$

For (ii): $\{u = \min(0, u_0(x^a)), x^a\}$

For (iii): The disjoint union of both.

It is now obvious that the Cauchy boundary is a C^0 manifold (actually locally Lipschitz) without boundary. The uniqueness statement follows from hypothesis A2. \square

5 Acknowledgments

M.M. is very grateful to Miguel Sánchez for valuable discussion on this subject. The authors wish to thank the Albert Einstein Institute and the University of Salamanca for their

hospitality and support. M.M. wishes to acknowledge financial support under the projects FIS2009-07238 (Spanish MICINN) and P09-FQM-4496 (Junta de Andalucía and FEDER funds).

6 Appendix A: Fixed points have $I_1 < 0$

In this Appendix we prove Lemma 2. Let p be a fixed point in $\partial^T \Sigma^T$. In fact, the lemma also holds for the more general case of $p \in \partial^T \{\lambda > 0\}$. For the purpose of the proof it is convenient to assume $p \in \partial^T \{\lambda > 0\}$ and extend the definition of horizon given in Definition 8 to any leaf \mathcal{L}_α intersecting $\partial^T \{\lambda > 0\}$. The constancy of the surface gravity and Proposition 6 also hold for such horizons.

We know that $I_1 \leq 0$ on $\{\lambda > 0\}$ (see (16)) and, by continuity $I_1(p) \leq 0$. So, we only need to exclude the possibility $I_1 = 0$. Let us assume that $I_1|_p = 0$ and find a contradiction.

Our aim is to show that there exists a non-degenerate horizon \mathcal{H}_α satisfying $p \in \overline{\mathcal{H}_\alpha}$. Since $I_1 = -2\kappa_\alpha^2$ (see Proposition 6) and κ_α is constant and non-zero on a non-degenerate horizon, we would contradict $I_1|_p = 0$. To that aim we only need to find a smooth path $\gamma(s)$ lying on a non-degenerate horizon and containing p in its closure.

First we note that $dY|_p \neq 0$ and $dN|_p \neq 0$. Indeed, if $dN|_p = 0$, then $Z|_p = 0$ and the definition of I_1 (together with $I_1 = 0$) implies $dY|_p = 0$. However, in Lemma 1 in [5] it is proved that a fixed point cannot have $dY = 0$ and $dN = 0$ unless the Killing data N, Y vanishes identically, which is not the case (the proof of Lemma 1 in [5] is done explicitly in dimension $m = 3$ but it carries through to arbitrary dimension with trivial changes). Thus, $dN|_p \neq 0$, and then the vanishing of $I_1|_p$ also implies $dY|_p \neq 0$.

Now, in Lemma 8 in [5] it is proved (again the proof is done there in dimension 3, but extends to arbitrary dimension) that there exists a positive constant b and a unit one-form $X \in T_p^* \Sigma$ orthogonal to $dN|_p$ such that

$$(34) \quad dY|_p = \frac{2b}{Q} (dN|_p \wedge X),$$

where $Q = |dN|_p|_g$. Evaluating I_1 at p we find

$$(35) \quad I_1|_p = 2(b^2 - Q^2).$$

Imposing $I_1|_p = 0$ we conclude $b = Q$. Let us now evaluate the Hessian of λ at p . Since $\lambda = N^2 - |Y|_g^2$ and p is a fixed point a simple calculation yields

$$\text{Hess}_g \lambda|_p = -2Q^2 X \otimes X.$$

By the Gromoll-Meyer splitting Lemma [17] there exist coordinates $\{y, x, z^A\}$ $A = 3, \dots, m$, in an open neighbourhood W_p of p such that λ takes the form $\lambda = -Q^2 y^2 + f(x, z^A)$ on

W_p for some function f and, moreover, p has coordinates $(0, \dots, 0)$, $X = dy|_p$, $dN|_p = dx|_p$ and f vanishes at p together with its gradient and its Hessian. From (34) we also have $Y = Q(xdy - ydx) + O(2)$. We are now in a position where the path $\gamma(s)$ mentioned above can be constructed.

For that we need to investigate the region $\{\lambda > 0\}$ near p . This region corresponds to $f > Q^2 y^2$. Since $p \in \partial^T \{\lambda > 0\}$ it is clear that $p \in \overline{\{f > 0\} \cap \{y = 0\}}$. If there exists a smooth curve $\tilde{\Gamma}(s) \subset \{y = 0\}$ approaching p and satisfying $f(\tilde{\Gamma}(s)) > 0$ then we are done because the curve $\Gamma(s) := \{y = Q^{-1} \sqrt{f(\tilde{\Gamma}(s))}, f(\tilde{\Gamma}(s))\}$ has the desired properties because (i) it lies on $\partial^T \{\lambda > 0\}$ and (ii) $d\lambda$ is nowhere zero on the curve (because $y \neq 0$ there) and hence $\Gamma(s)$ lies on a non-degenerate horizon.

So, it only remains to show that the curve $\tilde{\Gamma}(s)$ exists. Define $\mathcal{V} \subset \{y = 0\}$ as the set of points where the component Y_y does not vanish. Since $Y_y = Qx + O(2)$ it is clear that \mathcal{V} intersects $\{f > 0\}$ and also that there exists a smooth curve $\tilde{\Gamma}(s)$ fully contained in \mathcal{V} which approaches p . Our last step is to show that in fact $\mathcal{V} \subset \{f > 0\}$. Consider a smooth curve $\hat{\Gamma}(s)$ starting on a point $q \in \mathcal{V} \cap \{f > 0\}$. As long as f remains positive on this curve, consider the smooth curve $\bar{\Gamma}(s) := \{y = Q^{-1} \sqrt{f(\hat{\Gamma}(s))}, \hat{\Gamma}(s)\}$ which lies on a non-degenerate horizon. The y component of the equation $d\lambda = 2\kappa_\alpha Y$ on $\bar{\Gamma}(s)$ reads

$$-Q \sqrt{f(\hat{\Gamma}(s))} = \kappa_\alpha Y_y$$

Since κ_α is constant and $Y_y \neq 0$ on \mathcal{V} it follows that $f(\hat{\Gamma}(s))$ cannot become zero while remaining inside \mathcal{V} . This implies $\mathcal{V} \subset \{f > 0\}$ as claimed, and the lemma is proved. \square

References

- [1] R. Beig and P.T. Chruściel. Killing vectors in asymptotically flat space-times: I. asymptotically translational killing vectors and the rigid positive energy theorem. *J. Math. Phys.*, 37:1939–1961, 1996.
- [2] G. Bunting and A.K.M. Masood ul Alam. Nonexistence of multiple black holes in asymptotically euclidean static vacuum space-time. *Gen. Rel. Grav.*, 19:147–154, 1987.
- [3] C. Camacho and A.L. Neto. *Geometric theory of foliations*. Birkhauser, Boston Inc., 1985.
- [4] A. Carrasco. Trapped surfaces in spacetimes with symmetries and applications to uniqueness theorems. Ph.D. Thesis (2011).
- [5] A. Carrasco and M. Mars. On marginally outer trapped surfaces in stationary and static spacetimes. *Class Quantum Grav.*, 25:055011, 2008.

- [6] A. Carrasco and M. Mars. Uniqueness theorem for static spacetimes containing marginally outer trapped surfaces. *Class Quantum Grav.*, 28:175018, 2011.
- [7] I. Chavel. *Riemannian geometry, a modern introduction*. Cambridge Studies in Advanced Mathematics **98**, Cambridge University Press, 2006.
- [8] P.T. Chruściel and G.J. Galloway. Uniqueness of static black-holes without analyticity. *Class Quantum Grav.*, 27:152001, 2010.
- [9] P.T. Chruściel. The classification of static vacuum space-times containing an asymptotically flat spacelike hypersurface with compact interior. arXiv:gr-qc/9809088v2 (corrigendum to arXiv:gr-qc/9809088 submitted in 2010).
- [10] P.T. Chruściel. 'No hair' theorems - folklore, conjectures, results. *Contemporary Math.*, 170:23–49, 1994.
- [11] P.T. Chruściel. The classification of static vacuum space-times containing an asymptotically flat spacelike hypersurface with compact interior. *Class Quantum Grav.*, 16:661–687, 1999.
- [12] P.T. Chruściel and J. Lopes Costa. On uniqueness of stationary black holes. *Astérisque*, 321:195–265, 2008.
- [13] P.T. Chruściel and D. Maerten. Killing vectors in asymptotically flat space-times: II. asymptotically translational killing vectors and the rigid positive energy theorem in higher dimensions. *J. Math. Phys.*, 47:022502, 2006.
- [14] P.T. Chruściel and K.P. Tod. The classification of static electro-vacuum spacetimes containing an asymptotically flat spacelike hypersurface with a compact interior. *Commun. Math. Phys.*, 271:577–589, 2007.
- [15] T. Damour and B. Schmidt. Reliability of perturbation theory in general relativity. *J. Math. Phys.*, 31:2441–2453, 1990.
- [16] L.C. Evans and R.F. Gariepy. *Measure theory and fine properties of functions*. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1992.
- [17] D. Gromoll and W. Meyer. On differentiable functions with isolated critical points. *Topology*, 8:361–369, 1969.
- [18] S.W. Hawking and G.F.R. Ellis. *The large scale structure of space-time*. (Cambridge monographs on mathematical physics), Cambridge University Press, 1973.
- [19] C. Mantegazza and A.C Mennucci. Hamilton-jacobi equations and distance functions on riemannian manifolds. *Appl. Math. Optim.*, 47:1–25, 2003.

- [20] P. Miao. A remark on boundary effects in static vacuum initial data sets. *Class Quantum Grav.*, 22:L53–L59, 2005.
- [21] V. Moncrief. Spacetime symmetries and linearization stability of the einstein equations. *J. Math. Phys.*, 16:493–498, 1975.
- [22] R. Penrose. Gravitational collapse – the role of general relativity. *Nuovo Ciminto*, 1:252–276, 1965.
- [23] I. Rácz and R.M. Wald. Extensions of spacetimes with killing horizons. *Class Quantum Grav.*, 9:2643–2656, 1992.
- [24] A.D. Rendall. *Partial Differential equations in General Relativity*. Oxford Graduate Texts in Mathematics **16**, 2008.
- [25] R.M. Wald. *General Relativity*. The University of Chicago Press, 1984.
- [26] H. Weyl. Zur gravitationstheorie. *Ann. Phys. (Berlin)*, 54:117–145, 1917.